arXiv:1005.3512v2 [math.GT] 26 May 2010

A complete list of lens spaces constructed by Dehn surgery I

Motoo Tange

Abstract

Berge in [1] defined doubly primitive knots, which yield lens spaces by Dehn surgery. At the same paper he listed the knots into several types. In this paper we will prove the list is complete when $\tau > 1$. The invariant τ is a quantity with regard to lens space surgery, which is defined in this paper. Furthermore at the same time we will also prove that Table 6 in [8] is complete as Poincaré homology sphere surgery when $\tau > 1$.

1 Introduction

1.1 Several necessary conditions for lens space surgery

Let K be a knot in a 3-manifold M. Removing the open tubular neighborhood $\mathrm{nbd}(K)$ of K, and gluing the solid torus V by some map between the boundary of V and $M - \mathrm{nbd}(K)$ we obtain a new 3-manifold M'. By iterating some Dehn surgeries we can obtain all 3-manifolds. However we can not obtain all 3-manifolds by single Dehn surgeries of knots in S^3. When a lens space is given by a Dehn surgery of a knot, certain restrictions are imposed on the lens space and the knot. In this section we share with us some of the restrictions.

Let K be a knot in a homology sphere Y. The manifold $Y_p(K)$ stands for Dehn surgery of K with slope p. We define lens space $L(p,q)$ to be $S^3_{p/q}(\text{unknot})$. The problem of when $S^3_p(K)$ is lens space and when lens space $L(p,q)$ is obtained from Dehn surgery of K is fundamental.

Kadokami and Yamada proved the following by using torsion invariant.

Theorem 1 ([4]) *Let Y be a homology sphere. If $K \subset Y$ is a knot and $Y_p(K)$ is lens space $L(p,q)$, then the Alexander polynomial is the following form:*

$$\Delta_K(t) = \frac{(t^{hg}-1)(t-1)}{(t^g-1)(t^h-1)} \quad (t^p - 1)$$

up to multiplication of $\pm t^{\pm 1}$, where h, g are coprime integers and satisfy $hg = \pm 1 (p)$, and $h^2 = \pm q \ (p)$.

We can see that in the case where there exist h and g with $\gcd(h,g) = 1$ and $hg+1 \leq p$, the Alexander polynomial is the same as the polynomial of (h,g)-torus knot $T_{h,g}$. This type corresponds to type (I),(II) in Table 1. We have to notice that any coefficient in Theorem 1 is equivalent to the formula by using correction term in [6].

Ozsváth and Szabó [6], and Greene [3] showed that the Seifert genus $g(K)$ of any knot K yielding lens spaces has an inequality $2g(K) - 1 \leq p$ (Ozsváth-Szabó) and $2g(K) - 1 \leq p - 2\sqrt{\frac{4p+1}{5}}$ (Greene), where p is the order of H_1 of the lens space. The similar inequality holds for $\Sigma(2,3,5)$ (see [8]).

Let Y be a homology sphere. A Dehn surgery $Y_p(K)$ has the core circle of the surgery which is called the circle *dual knot*. In the case where $Y_p(K)$ is lens space $L(p,q)$, the first homology class of the dual knot $\tilde{K}$ assigns an integer k. In fact since there is a core circle by genus 1 Heegaard splitting of the lens space, we have only to take k as the difference $[\tilde{K}]=k[c]$, where c is the core circle so that $q=k^2(p)$ holds. However the assignment has two ambiguities: the choice of the non-trivial core circle in the two solid tori and the choice of the orientation of the knot. Thus we consider the integer k as the set $\mathcal{D}(p,K):=\{k,-k,k^{-1},-k^{-1}\}\subset\mathbb{Z}/p\mathbb{Z}$. Here we call it *the dual class invariant.* The integers h,g in Theorem 1 are elements of $\mathcal{D}(p,K)$. Abstractly we define $\mathcal{D}(p,k)$ to be the set $\{k,-k,k^{-1},-k^{-1}\}$.

Here we define a Laurent polynomial as follows. Let $T_{r,s}$ be the (r,s)-torus knot.

Definition 1 *Let (p,k) be a coprime integers pair, and h,g two coprime representations in $\mathcal{D}(p,k)$ with $hg=\pm1$ (p). Adding $\pm t^n(t^p-1)$ several times to the symmetrized Alexander polynomial $\Delta_{T_{h,g}}(t)$, we can make any exponent of the terms of the polynomial change a term between $-\lfloor\frac{p}{2}\rfloor+1$ and $\lfloor\frac{p}{2}\rfloor$. Then we get a Laurent polynomial $\tilde{\Delta}_{p,k}(t)$. If p is even and the $p/2$-th term $a_{\frac{p}{2}}$ is not zero, then we modify $\tilde{\Delta}_{p,k}(t)$ into*

$$\tilde{\Delta}_{p,k}(t)+\frac{a_{\frac{p}{2}}}{2}t^{-\frac{p}{2}}-\frac{a_{\frac{p}{2}}}{2}t^{\frac{p}{2}}$$

In this way we get a symmetric Laurent polynomial and denote it by $\Delta_{p,k}(t)$.

We define $g(p,k_1)$ to be the degree of $\Delta_{p,k_1}(t)$.

By Fintushel and Stern's work in [2] conversely if there exists a coprime integer pair (p,k) satisfying $q=k^2(p)$, one can realize a Dehn surgery $L(p,q)=Y_p(K)$ on a homology sphere Y with $k\in\mathcal{D}(p,K)$. Hence one of lens space surgery problems is to consider when for a coprime pair (p,k) there exists a knot K in S^3 (in general in a fixed homology sphere) with $k\in\mathcal{D}(p,K)$.

Our starting point is a coprime pair (p,k) then we call (p,k) *the initial data* for lens space surgery by taking the minimal k in $\mathcal{D}(p,k)$. Here any element in $\mathcal{D}(p,k)$ is reduced between 0 and $p-1$. For an initial data (p,k_1) if there exists a lens space surgery on a homology sphere Y such that $k\in\mathcal{D}(p,K)$, where K is the knot in Y, we call (p,k) the initial data *realized by the lens space surgery.*

We introduce an explicit coefficient formula of $\Delta_{p,k}(t)$, which is computed by Theorem 1, that was proved in [7]. Let (p,k) be a coprime integer satisfying $0<k<p$ and k' the inverse of $k \mod p$ satisfying $0<k'<p$. We use this notation in other places as long as we do not indicate. We put $m:=\frac{kk'-1}{p}$, $q=k^2(p)$, $c:=\frac{(k+1-p)(k-1)}{2}$, and

$$\Phi^l_{p,q}(k)=\#\{j\in\{1,2,\cdots k'\}|[qj-l]_p\in\{1,2,\cdots,k\}\},$$

where $[\alpha]_p$ stands for the reduction of any integer α to $0\le[\alpha]_p<p$.

Theorem 2 ([8]) *Let (p,k) be a coprime integer pair and a_i the i-th coefficient of the Laurent polynomial $\tilde{\Delta}_{p,k}(t)$. Then a_i $(-\lfloor\frac{p}{2}\rfloor+1\le i\le\lfloor\frac{p}{2}\rfloor)$ has the following explicit formula*

$$a_i=-m+\Phi^{ki+c}_{p,q}(k). \tag{1}$$

Hence if a knot $K\subset S^3$ yields lens space $L(p,q)$, the symmetrized Alexander polynomial $a_i(K)$ is computed by the formula. From the genus bound $2g(K)-1\le p$ by Ozsváth and Szabó we can get the following.

Corollary 1 *For a lens space surgery $L(p,q)=S^3_p(K)$ with initial data (p,k) we have*

$$\Delta_K(t)=\Delta_{p,k}(t).$$

In particular $g(K)=g(p,k_1)$ holds.

When we replace S^3 with $\Sigma(2,3,5)$, we get the same equality. In this paper we assume that the genus of any knot yielding lens space surgery satisfies $2g(K) \leq p$. Ozsváth and Szabó's result does not deny the $2g(K)-1=p$ case. The author was informed that Dr. Greene showed that at least the $2g(K)-1=p$ case can be ruled out as long as lens space surgery over S^3.

Ozsváth and Szabó proved the new restrictions in [5, 6].

Theorem 3 ([5]) *If $K \subset S^3$ is a knot and $S^3_p(K)$ is lens space, then the symmetrized Alexander polynomial $\Delta_K(t)$ is the following form:*

$$\Delta_K(t) = (-1)^m + \sum_{j=1}^{m}(-1)^k(t^{n_j}+t^{-n_j}),$$

where the sequence $\{n_j\}$ is increasing natural numbers: $n_1 < n_2 < \cdots < n_m$.

The same assertion is also satisfied for $\Sigma(2,3,5)$ in place of S^3.

Theorem 4 ([6]) *If $K \subset S^3$ is a knot and $S^3_p(K)$ is a lens space $L(p,q)$, then the following quantity*

$$t_i := \begin{cases} d(L(p,q),ki+c) - d(L(p,1),i) & \text{if } 2|i| \leq p \\ 0 & \text{otherwise.} \end{cases}$$

is non-negative integer and is coincident with the torsion invariant $t_i(K)$ for any integer i. Here $t_i(K)$ is the i-th Turaev torsion of $S^3_0(K)$.

Some of lens space surgeries on $\Sigma(2,3,5)$ fail to this theorem, for example $L(22,3)$ is obtained as a lens space surgery on $\Sigma(2,3,5)$ and $t_{11}=-2$ holds.

By using these restrictions, we can make a sharp distinction as in [5]. However either of Theorem 3 and 4 is not perfect so as to distinguish lens space surgery.

1.2 Berge's examples

Berge in [1] defined a class of knots in S^3, which is most important to lens space surgery so far.

Definition 2 *Let K be a knot in S^3. We call K a doubly primitive knot if K isotopic to a knot L in a genus 2 Heegaard surface in S^3 and both classes induced in $\pi_1(V_i)$ are primitive elements, where V_i $(i=1,2)$ are the genus 2 handlebodies.*

This definition can be easily extended to any knot in a homology sphere with genus 2 Heegaard decomposition, which satisfies the same conditions.

He proved that all doubly primitive knots yield lens spaces by Dehn surgery with an integer slope. In the other words this condition is a sufficient condition for lens space surgery. He conjectures that doubly primitive knots are all knots yielding lens spaces. This conjecture is still open.

He also listed the doubly primitive knots in [1]. Thus any of the list is realized by a lens space surgery on S^3, however it is open question that this list is complete.

In [9] J. Rasmussen rearranged the list to the equivalent one below (Table 1).

1.3 Poincaré homology sphere version of Berge's examples

Next we introduce the result [8], which is a sufficient condition for lens space surgery on Poincaré homology sphere.

I,II	$p = ik \pm 1\ (k^2)$	$\gcd(i,k) = 1, 2$
III$_\pm$	$p = (2k \mp 1)J\ (k^2)$	$J \in \mathbb{Z}; J \vert k \pm 1; \frac{k \pm 1}{J}:$ odd
IV$_\pm$	$p = (k \pm 1)J\ (k^2)$	$J:$ odd; $J \vert 2k \mp 1$
V$_\pm$	$p = (k \pm 1)J\ (k^2)$	$J \in \mathbb{Z}; J \vert k \pm 1; J:$ odd
VII, VIII	$k^2 \pm k \pm 1 = 0(p)$	
IX	$p = 22J^2 + 9J + 1$	$k = 11J + 2, J \in \mathbb{Z}$
X	$p = 22J + 13J + 2$	$k = 11J + 3, J \in \mathbb{Z}$

Table 1: Berge's list

type	p	k
A$_1$	$14J^2 + +7J + 1$	$7J + 2$
A$_2$	$20J^2 + 15J + 3$	$5J + 2$
B	$30J^2 + 9J + 1$	$6J + 1$
C$_1$	$42J^2 + 23J + 3$	$7J + 2$
C$_2$	$42J^2 + 47J + 13$	$7J + 4$
D$_1$	$52J^2 + 15J + 1$	$13J + 2$
D$_2$	$52J^2 + 63J + 19$	$13J + 8$
E$_1$	$54J^2 + 15J + 1$	$27J + 4$
E$_2$	$54J^2 + 39J + 7$	$27J + 10$
F$_1$	$69J^2 + 17J + 1$	$23J + 3$
F$_2$	$69J^2 + 29J + 3$	$23J + 5$
G$_1$	$85J^2 + 19J + 1$	$17J + 2$
G$_2$	$85J^2 + 49J + 7$	$17J + 5$
H$_1$	$99J^2 + 35J + 3$	$11J + 2$
H$_2$	$99J^2 + 53J + 7$	$11J + 3$
I$_1$	$120J^2 + 16J + 1$	$12J + 1$
I$_2$	$120J^2 + 20J + 1$	$20J + 2$
I$_3$	$120J^2 + 36J + 3$	$12J + 2$
J	$120J^2 + 104J + 22$	$12J + 5$
K	191	15

Table 2: The Poincaré homology sphere version of Berge's examples.

Theorem 5 ([8]) *Let (p,k) be any of Table 2. Then (p,k) is realized by a lens space surgery on $K \subset \Sigma(2,3,5)$. Furthermore we can take a doubly primitive knot as a knot for the surgery.*

Theorem 1 and Theorem 3 also hold for lens space surgery on the Poincaré homology sphere. However Theorem 4 does not hold because the initial data $(22,5)$ does not satisfy Theorem 4 but admits lens space surgery $\Sigma(2,3,5)_{22}(K)$.

Then the following is conjectured.

Conjecture 1 *Let (p,k) be a coprime integer. Suppose that the polynomial $\Delta_{k,p}(t)$ satisfies the alternating condition in Theorem 3, and the pair satisfies the condition in Theorem 4. Then (p,k) is realized by a lens space surgery on S^3.*

This conjecture is equivalent to Conjecture 1.12 in [6].

Conjecture 2 *Let (p,k) be a coprime integer. Suppose that the polynomial $\Delta_{p,k}(t)$ satisfies the alternating condition in Theorem 3, Then (p,k) is realized by a lens space surgery on S^3 or $\Sigma(2,3,5)$.*

If Conjecture 1 is true, then it means that Table 1 is the complete list of doubly primitive knots. If Conjecture 2 is true, then it means that Table 1 and 2 are the complete list of doubly primitive knots in S^3 and $\Sigma(2,3,5)$. Even if these conjectures are proven, the problem of whether doubly primitive knots are all the knots yielding lens space or not remains open. The aim of this paper is a partial contribution to Conjecture 1 and 2. The meaning of "partiality" is described later.

2 Preliminaries

2.1 Quadratic relations for lens space surgery

In this section we shall define a quadratic relation in $\mathbb{Z}/p\mathbb{Z}$. Let (p,k_1) be a coprime integer pair with p positive and k_2 the reduced element with $k_1 \leq k_2 < \frac{p}{2}$ in $\{k_1, [-k_1]_p, [k_1^{-1} \bmod p]_p, [-k_1^{-1} \bmod p]_p\}$.

We denote by $[[\alpha]]_p$ the reduction between $-\lfloor \frac{p}{2} \rfloor + 1$ and $\lfloor \frac{p}{2} \rfloor$. We define q_2 to be $[[k_2^2]]_p$ and a to be $||q_2| - k_2|$. Hence we can find a relation on $\mathbb{Z}/p\mathbb{Z}$

$$ak_1^2 + \epsilon_1 k_1 + \epsilon_2 = 0 \ (p), \tag{2}$$

where $\epsilon_i = \pm 1$. This relation is called *the associated (quadratic) relation* in this paper. Note that relations of (p,k_1) in the form of (2) are not always unique for any data. For example $(43,12)$ has two relations $2k_1^2 + k_1 + 1 = 0 \ (43)$ and $5k_1^2 + k_1 - 1 = 0 \ (43)$.

Proposition 1 *The $a = 0$ case is equivalent to $k_1 = k_2 = 1$ then this initial data can be realized by p-surgery of the unknot.*

Proof. Therefore $k_1 = \pm 1 \ (p)$ holds. □

The $a = 1$ case is (VII) and (VIII) in Table 1. Then we assume that $a \geq 2$.

Lemma 1 *Let (p,k_1) be an initial data with $k_1 > 1$. The associated quadratic relation of (p,k_1) is the relation which the term a, which is the coefficient of degree 2 of the relation, is minimal among the relations having form of (2).*

Any relation of the form of (2) with the minimal degree 2 term is the associated relation.

Proof The former of the assertion is obvious from the definition of a. We show the latter part. Suppose that (p,k_1) satisfies $ak_1^2+\epsilon_1k_1+\epsilon_2=0\ (p)$ and $ak_1^2+\epsilon_1'k_1+\epsilon_2'=0\ (p)$, and $\epsilon_1<\epsilon_1'$. Then we have $(\epsilon_1'-\epsilon_1)k_1=\epsilon_2-\epsilon_2'\ (p)$. Since we have $0<2k_1<p$, $2k_1=2$ holds. Then we have $k_1=1$. If $\epsilon_1=\epsilon_1'$, then this relation is the same. $\square$

Now let (p,k_1) be a coprime pair as above. We take a relation $ak_1^2+\epsilon_1k_1+\epsilon_2=0\ (p)$ and define an integer n to be $ak_1^2+\epsilon_1k_1+\epsilon_2-np=0$. Solving the quadratic equation, we get

$$k_1=\frac{-\epsilon_1+X}{2a}$$

where X is positive integer and we put $X^2=1-4a(\epsilon_2-np)$. Hence we have $p=\frac{X^2-D}{4an}$, where we put $D=1-4a\epsilon_2$. Here let τ,γ be integers satisfying $X=2an\tau+\gamma$ and $0\le\gamma<2an$. Hence we have

$$k_1=n\tau+\frac{\gamma-\epsilon_1}{2a}=n\tau+\gamma', \tag{3}$$

where we put $\gamma'=\frac{\gamma-\epsilon_1}{2a}$ $(-\frac{\epsilon_1}{2a}\le\gamma'<n-\frac{\epsilon_1}{2a})$, thus $\gamma=\epsilon_1\ (2a)$ holds. Then p is described as follows:

$$p=an\tau^2+\gamma\tau+\frac{\gamma^2-D}{4an}. \tag{4}$$

Since $\gamma^2-4an\frac{\gamma^2-D}{4an}=D=1-4a\epsilon_2$, if $\epsilon_2=1$, then the quadratic function has a positive value for any τ as long as $a>0$. If $\epsilon_2=-1$, then considering $p=an\left(\tau+\frac{\gamma}{2an}\right)^2-\frac{D}{4an}$, we have $an\left(0+\frac{\gamma}{2an}\right)^2-\frac{D}{4an}=\frac{a(\gamma')^2+\epsilon_1\gamma'-1}{n}\ge 0$ as long as $a>0$ and $\gamma'>1$. Thus we have $an\left(-2+\frac{\gamma}{2an}\right)^2-\frac{D}{4an}=4an-2\gamma+\frac{a(\gamma')^2+\epsilon_1\gamma'-1}{n}\ge 0$

Definition 3 *Let (p,k_1) be an initial data. We denote by τ the parameter computed as above by using the associated quadratic relation. If the parameter $\tau\ge 2$ holds we call (p,k_1) a stable (initial) data.*

Here we list the associated relation for lens surgeries in Table 1 and (2), except (I), (II), ($\text{V}_\pm$), (VII), and (VIII).

Here we state Main theorem in this paper.

Theorem 6 (Main theorem) *Suppose that (p,k_1) is a stable data. If $\Delta_{p,k}(t)$ satisfies the alternating condition in Theorem 1 and Theorem 3, then (p,k) is realized by a lens space surgery of types below:*

$$I, II, III_\pm, IV_\pm, VII, VIII, IX, X, A_i, C_i, D_i, E_i, F_i, G_i, H_i,$$

where $i=1$ or 2.

The partiality in Section 1 means the stability.

Lemma 2 *If $\tau\ge 2$ then, we have $k_2=ak_1+\epsilon_1$.*

Proof. Putting $\kappa=ak_1+\epsilon_1$, we have $p=\kappa(\tau+\frac{\gamma'}{n})+\frac{\epsilon_2}{n}$ and

$$-\frac{\epsilon_1}{2an}<\frac{\gamma'}{n}<1-\frac{\epsilon_1}{2an}.$$

Hence $\kappa\le\frac{p}{2}$ and $\kappa k_1=\pm 1\ (p)$, therefore we have $\kappa=k_2$. $\square$

In this section we shall concentrate on seeing the coefficients of the polynomial $\tilde{\Delta}_{p,k_1}(t)$ for the stable data (p,k_1).

Table 4 is the list of lens spaces with $\tau=1$ in Table 2 in [8]. These lens spaces are not dealt with in this paper. We conjecture that all lens spaces obtained by Dehn surgeries on $\Sigma(2,3,5)$ with $\tau=1$ are included in Table 4.

type	J	the associated relation	n
$\mathrm{III}_\pm$	$\lvert J\rvert \geq 2$	$ak_1^2 \pm k_1 - 1 = 0 \ \ n \mid a-2$	
$\mathrm{IV}_\pm$	$\lvert J\rvert \geq 2$	$ak_1^2 \pm k_1 - 1 = 0 \ \ n \mid a-2$	
IX,X	$\lvert J\rvert \geq 2$	$2k_1^2 \pm k_1 + 1 = 0$	11
A_1	$\lvert J\rvert \geq 2$	$2k_1^2 - \delta k_1 + 1 = 0$	7
A_2	$\lvert J\rvert \geq 2, J = 1$	$4k_1^2 - \delta k_1 + 1 = 0$	5
$\mathrm{E}_1, \mathrm{E}_2$	$\lvert J\rvert \geq 2$	$2k_1^2 - \delta k_1 - 1 = 0$	27
$\mathrm{F}_1, \mathrm{F}_2$	$\lvert J\rvert \geq 2$	$3k_1^2 - \delta k_1 - 1 = 0$	23
D_1	$\lvert J\rvert \geq 2$	$4k_1^2 - \delta k_1 - 1 = 0$	13
D_2	$\lvert J\rvert \geq 2, J = 1$	$4k_1^2 - \delta k_1 - 1 = 0$	13
$\mathrm{G}_1, \mathrm{G}_2$	$\lvert J\rvert \geq 2, J = 1$	$5k_1^2 - \delta k_1 - 1 = 0$	17
$\mathrm{C}_1, \mathrm{C}_2$	$\lvert J\rvert \geq 2, J = 1$	$6k_1^2 - \delta k_1 - 1 = 0$	7
$\mathrm{H}_1, \mathrm{H}_2$	$\lvert J\rvert \geq 2, J = 1$	$9k_1^2 - \delta k_1 - 1 = 0$	11
B	$J \geq 2$	$(15J^2 - 18J - 5)k_1^2 - k_1 - 1 = 0$	$18J^2 - 21J - 7$
B	$J \leq -2$	$(15J^2 + 27J + 6)k_1^2 - k_1 + 1 = 0$	$18J^2 + 33J + 8$
B	$J = 1$	$8k_1^2 + k_1 + 1 = 0$	10
I_1	$J \geq 2$	$(40J^2 - 28J - 3)k_1^2 - k_1 - 1 = 0$	$48J^2 - 32J - 5$
I_1	$J \leq -1$	$(40J^2 + 12J - 1)k_1^2 - k_1 - 1 = 0$	$48J^2 + 16J - 1$
I_2	$J \geq 2$	$(60J^2 - 50J - 5)k_1^2 - k_1 - 1 = 0$	$200J^2 - 160J - 23$
I_2	$J \leq -2$	$(60J^2 + 70J + 6)k_1^2 - k_1 + 1 = 0$	$200J^2 + 240J + 27$
I_3	$J \geq 2$	$(60J^2 - 42J - 9)k_1^2 - k_1 - 1 = 0$	$72J^2 - 48J - 13$
I_3	$J \leq -2$	$(60J^2 + 78J + 12)k_1^2 - k_1 + 1 = 0$	$72J^2 + 96J + 17$
J	$J \geq 1$	$(40J^2 + 28J + 6)k_1^2 + k_1 - 1 = 0$	$48J^2 + 32J + 7$
J	$J \leq -2$	$(40J^2 - 12J - 4)k_1^2 + k_1 - 1 = 0$	$48J^2 + 80J + 27$
K		$22k_1^2 + k_1 + 1 = 0$	26

Table 3: The associated relations. ($\delta = \mathrm{sgn}(J)$).

(p, k_1)	(type, J)	the associated relation
$(43, 12)$	$(\mathrm{A}_1, -2), (\mathrm{F}_2, -1), (\mathrm{G}_2, -1)$	$2k_1^2 + k_1 + 1 = 0$
$(8, 3)$	$(\mathrm{A}_1, -1), (\mathrm{A}_2, -1), (\mathrm{C}_2, -1), (\mathrm{D}_2, -1)$	$2k_1^2 - k_1 + 1 = 0$
$(38, 7)$	$(\mathrm{A}_2, 1), (\mathrm{D}_1, -1), (\mathrm{J}, -1)$	$4k_1^2 - k_1 + 1 = 0$
$(53, 8)$	$(\mathrm{A}_2, -2), (\mathrm{F}_1, -1), (\mathrm{H}_2, -1)$	$4k_1^2 + k_1 + 1 = 0$
$(68, 9)$	$(\mathrm{C}_1, 1), (\mathrm{D}_1, 1)$	$6k_1^2 - k_1 - 1 = 0$
$(125, 12)$	$(\mathrm{C}_1, -2)$	$6k_1^2 + k_1 - 1 = 0$
$(102, 11)$	$(\mathrm{C}_2, 1)$	$6k_1^2 - k_1 - 1 = 0$
$(87, 10)$	$(\mathrm{C}_2, -2), (\mathrm{I}_2, -1), (\mathrm{F}_1, 1)$	$6k_1^2 + k_1 - 1 = 0$
$(179, 24)$	$(\mathrm{D}_1, -2)$	$4k_1^2 + k_1 - 1 = 0$
$(134, 21)$	$(\mathrm{D}_2, 1)$	$4k_1^2 - k_1 - 1 = 0$
$(101, 18)$	$(\mathrm{D}_2, -2), (\mathrm{I}_2, -1), (\mathrm{F}_2, 1)$	$4k_1^2 + k_1 - 1 = 0$
$(187, 50)$	$(\mathrm{E}_1, -2)$	$2k_1^2 + k_1 - 1 = 0$
$(145, 44)$	$(\mathrm{E}_2, -2)$	$2k_1^2 + k_1 - 1 = 0$
$(243, 43)$	$(\mathrm{F}_1, -2)$	$3k_1^2 + k_1 - 1 = 0$
$(221, 41)$	$(\mathrm{F}_2, -2)$	$3k_1^2 + k_1 - 1 = 0$
$(303, 32)$	$(\mathrm{G}_1, -2)$	$5k_1^2 + k_1 - 1 = 0$
$(249, 29)$	$(\mathrm{G}_2, -2)$	$5k_1^2 + k_1 - 1 = 0$
$(141, 22)$	$(\mathrm{G}_2, 1), (\mathrm{I}_2, 1)$	$5k_1^2 - k_1 - 1 = 0$
$(329, 20)$	$(\mathrm{H}_1, -2)$	$9k_1^2 + k_1 - 1 = 0$
$(137, 13)$	$(\mathrm{H}_1, 1), (\mathrm{I}_1, 1)$	$9k_1^2 - k_1 - 1 = 0$
$(297, 19)$	$(\mathrm{H}_2, -2)$	$9k_1^2 + k_1 - 1 = 0$
$(159, 14)$	$(\mathrm{H}_2, 1), (\mathrm{I}_2, 1)$	$9k_1^2 - k_1 - 1 = 0$

Table 4: The associated relations having lens space surgeries on $\Sigma(2,3,5)$ with $\tau = 1$.

2.2 The $\gamma' = 0$ or $\gamma' = n$ case.

First we classify all stable data realized lens space surgery on S^3 or $\Sigma(2,3,5)$ in the cases of $(\epsilon_1, \gamma') = (1,0), (-1,n)$.

Theorem 7 *In the case of* $(\epsilon_1, \gamma') = (1,0)$ *the data is realized by type (I).*

Proof. In the case of $(\epsilon_1, \gamma') = (1,0)$ we have $\gamma = 1$. From integrality of p and Formula (4), $\frac{1}{n}$ is an integer thus $n = 1$. Therefore $(p, k_1) = (a\tau^2 + \tau + \epsilon_2, \tau)$ holds. Thus we have $k_2 = a\tau + 1$. From this and Theorem 12 in [7], $\gcd(k_1, k_2) = 1$ or 2. Therefore this data is realized by type (I).

In the case of $(\epsilon_1, \gamma') = (-1, n)$ we have $\gamma = 2an - 1$. In the same way from integrality of p, we get $n = 1$. Thus $(p, k) = ((\tau+1)(a(\tau+1)-1) + \epsilon_2, \tau+1)$. Hence we have $k_2 = a(\tau+1) - 1$. From Theorem 12 in [7], $\gcd(k_1, k_2) = 1$ or 2. This data is realized by type (I). $\square$

Then the inverse element k_1' mod p is reduced as follows:

$$k_1' := \begin{cases} k_2 & \epsilon_2 = -1 \\ p - k_2 & \epsilon_2 = 1. \end{cases}$$

2.3 The case of $k_1 = 1, 2, 3$.

In the case of $k_1 = 1$, k_2 is also 1 and the initial data is realized by the unknot surgery $L(p,1) = S^3_p(\text{unknot})$.

In the case of $k_1 = 2$, p is odd. Then we put p by $2n+1$. Hence $k_2 = n$ holds. From this and Theorem 12 in [7], n is odd. In this case each of data is realized by the $(2,n)$-torus knot surgery.

In the case of $k_1 = 3$, p is $3n \pm 1$. Then $k_2 = n$ holds. From this and Theorem 12 in [7], $\gcd(3,n) = 1$ holds. In this case each of data is realized by the $(3,n)$-torus knot surgery.

Therefore now we may assume that $k_1 \geq 4$ holds.

2.4 The $n = 1$ case.

If $n = 1$ holds, then we have $ak_1^2 + \epsilon_1 k_1 + \epsilon_2 = p$. This data is realized by $(k_1, ak_1 + \epsilon_1)$-torus knot surgery. This is type (I) in Berge's list.

Therefore we assume that $n \geq 2$.

2.5 The coefficients of the Alexander polynomial

For a stable data (p, k_1) the reduced square q_2 satisfies $q_2 = \epsilon_1 k_2 - \epsilon_2 a$ and we have $k_1' = [-\epsilon_2 k_2]_p$. We put $q_1 := [k_1^2]_p$, $c := \frac{(k_1+1-p)(k_1-1)}{2}$ and $\bar{g} = k_1'(c - q_1) + 1 = -\epsilon_2 k_2 (c - q_1) + 1 \ (p)$. Then we get

$$-2\bar{g} = k_1 + k_1' - 2 = k_1 - \epsilon_2 k_2 - 2 \ (p) \tag{5}$$

We define a function A to be

$$A(n_1, n_2, n_3) = \tilde{a}_{-\bar{g} + n_1 k_1 + n_2 k_1' + n_3},$$

where the coefficients $\tilde{a}_i = a_{[[i]]_p}$ for any integer $i \in \mathbb{Z}$. The coefficients $\tilde{a}_i$ are regarded as the function that cyclically lifts the coefficients of $\Delta_{p,k_1}(t)$ to $\mathbb{Z}$.

Here we prepare the following lemma.

Lemma 3 *Let (p, k_1) be any initial data. We have*

$$A(m_1, m_2, m_3) - A(m_1, m_2 - 1, m_3) = E_{k_1'}(m_2q_2 + m_3k_1' + m_1 + 1) - E_{k_1'}(m_2q_2 + (m_3 - 1)k_1' + m_1 + 1)$$

$$= E_{p-k_1'}(m_2q_2 + (m_3 - 2)k_1' + m_1 + 1) - E_{p-k_1'}(m_2q_2 + (m_3 - 1)k_1' + m_1 + 1),$$

where the function $E_x(y)$ is defined to be

$$E_x(y) = \begin{cases} 1 & y \equiv 1, 2, \cdots, [x]_p \mod p \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. By definition we have

$$\Phi_{p,q_1}^{k_1 i + c}(k_1) = \sum_{j=1}^{k_1} E_{k_1'}(q_2 j + k_1' i + q_2 c).$$

Now we put $\sigma = -\bar{g} + m_1 k_1 + m_3$

$$\begin{aligned} & A(m_1, m_2, m_3) - A(m_1, m_2 - 1, m_3) \\ = & \sum_{j=1}^{k_1} E_{k_1'}(q_2 j + k_1'(\sigma + m_2 k_1') + q_2 c) - \sum_{j=1}^{k_1} E_{k_1'}(q_2 j + k_1'(\sigma + (m_2 - 1)k_1') + q_2 c) \\ = & \sum_{j=1}^{k_1} E_{k_1'}(q_2 j + k_1'(\sigma + m_2 k_1') + q_2 c) - \sum_{j=0}^{k_1 - 1} E_{k_1'}(q_2 j + k_1'(\sigma + m_2 k_1') + q_2 c) \\ = & E_{k_1'}(m_2 q_2 + (\sigma + 1)k_1' + q_2 c) - E_{k_1'}(m_2 q_2 + \sigma k_1' + q_2 c). \end{aligned} \tag{6}$$

Here we have

$$\begin{aligned} \sigma k_1' + q_2 c &= (-\bar{g} + m_1 k_1 + m_3)k_1' + q_2 c \\ &= -(k_1'(c - q_1) + 1)k_1' + m_1 + m_3 k_1' + q_2 c = (m_3 - 1)k_1' + m_1 + 1 \ (p). \end{aligned}$$

Therefore we have

$$(6) = E_{k_1'}(m_2 q_2 + m_3 k_1' + m_1 + 1) - E_{k_1'}(m_2 q_2 + (m_3 - 1)k_1' + m_1 + 1).$$

Furthermore, by using the obvious relation $E_{k_1'}(x) + E_{p-k_1'}(x - k_1') = 1$ we get

$$(6) = E_{p-k_1'}(m_2 q_2 + (m_3 - 2)k_1' + m_1 + 1) - E_{p-k_1'}(m_2 q_2 + (m_3 - 1)k_1' + m_1 + 1).$$

□

We can describe the formulas of Lemma 3 as follows:

$$\begin{aligned} & A(m_1, m_2, m_3) - A(m_1, m_2 - 1, m_3) \\ = & E_{k_2}(m_2 q_2 - \epsilon_2 m_3 k_2 + m_1 + 1) - E_{k_2}(m_2 q_2 - \epsilon_2 (m_3 - 1)k_2 + m_1 + 1). \end{aligned} \tag{7}$$

2.6 The two reductions: q_1 and q_2.

Let (p, k_1) be a stable data. Suppose that (p, k_1) satisfies a relation

$$a k_1^2 + \epsilon_1 k_1 + \epsilon_2 = 0. \ (p)$$

Here we recall $a = -\epsilon_2 q_2 + \epsilon_2 \epsilon_1 k_2$. Then we have

$$k_2 = \epsilon_1 q_2 + \epsilon_1 \epsilon_2 a. \tag{8}$$

Hence

$$q_2 = \epsilon_1 a k_1 - \epsilon_2 a + 1 \text{ and } |q_2| = \epsilon_1 q_2.$$

From

$$\frac{|q_2|}{a} = k_1 - \epsilon_1\epsilon_2 + \frac{\epsilon_1}{a}$$

we have $\frac{|q_2|}{a} \geq 4 - \epsilon_1\epsilon_2 + \frac{1}{a} > 2$, as a result $\frac{a}{|q_2|} < \frac{1}{2}$ holds. Furthermore from $\epsilon_1 q_2 = k_2 - \epsilon_1\epsilon_2 a \geq 4a + \epsilon_1 - \epsilon_1\epsilon_2 a \geq 3a - 1 > 0$,

$$|q_2| = \epsilon_1 q_2$$

Let η be the integer satisfying $aq_1 + \epsilon_1 k_1 + \epsilon_2 = \eta p$ and $0 < q_1 < p$.

$$q_1 = \frac{\eta p - \epsilon_1 k_1 - \epsilon_2}{a},$$

then $k_1^2 - q_1 = \frac{n-\eta}{a} p$ implies $\eta = n$ (a), namely $\eta = [n]_a$.

2.7 A symmetry of the stable data

Over all stable data $\{(p, k_1)\}$ we define an involution as follows:

$$(p, k_1) \to (p - (\gamma - an)(2\tau + 1), k_1 + n - 2\gamma') := (\underline{p}, \underline{k}_1)$$

$$(\gamma', \gamma, X) \to (n - \gamma', 2an - \gamma, X + 2(an - \gamma)$$

$$(\epsilon_1, \epsilon_2) \to (-\epsilon_1, \epsilon_2)$$

$$(a, n, \tau, D) \to (a, n, \tau, D).$$

In fact we have $\underline{k}_1 \geq k_1 - n = n(\tau - 1) + \gamma' \geq 0$ and $\underline{p} = an(\tau + 1)^2 - \gamma(\tau + 1) + \frac{\gamma^2 - D}{4an}$

$$\begin{aligned} a\underline{k}_1^2 - \epsilon_1\underline{k}_1 + \epsilon_2 &= ak_1^2 + \epsilon_1 k_1 + \epsilon_2 + 2ak_1(n - 2\gamma') + a(n - 2\gamma')^2 - \epsilon_1(n - 2\gamma') - 2\epsilon_1 k_1 \\ &= n(p + (an - \gamma)(2\tau + 1)) = n\underline{p}. \end{aligned}$$

Hence $\underline{p}$ and $\underline{k}_1$ are relatively prime and $\underline{p} > 0$. Similarly putting

$$\underline{k}_2 := a\underline{k}_1 - \epsilon_1 = ak_1 + an - 2a\gamma' - \epsilon_1 = k_2 + an - \gamma - \epsilon_1$$

we have $\frac{1}{n}(\underline{k}_1\underline{k}_2 + \epsilon_1) = \underline{k}_2(\tau + 1 - \frac{\gamma'}{n}) + \frac{\epsilon_1}{n} \geq 2\underline{k}_2 + \frac{\epsilon_1}{n}$, hence $\underline{k}_2 \leq \frac{1}{2}(\underline{p} - \frac{\epsilon_2}{n})$. Therefore $\{\underline{k}_1, \underline{k}_2, p - \underline{k}_2, p - \underline{k}_1\}$ is the dual class invariant for the data $(\underline{p}, \underline{k}_1)$ and $0 < \underline{k}_1 \leq \underline{k}_2 < \frac{\underline{p}}{2}$ is satisfied.

Lemma 4 *Let (p, k_1) be a stable data. Let $(\underline{p}, \underline{k}_1)$ be the data from (p, k_1) via the involution. Then $a\underline{k}_1^2 - \epsilon_1\underline{k}_1 + \epsilon_2 = 0(\underline{p})$ is the associated relation.*

Proof. Since $\underline{q}_2 = [[\underline{k}_2^2]]_{\underline{p}} = -\epsilon_1\underline{k}_2 - \epsilon_2 a$, the absolute value $|\underline{q}_2| = -\epsilon_1\underline{q}_2 = \underline{k}_2 + \epsilon_1\epsilon_2 a$. Hence we have $||\underline{q}_2| - k_2| = a$. Therefore $a\underline{k}_1^2 - \epsilon_1\underline{k}_1 + \epsilon_1 = 0(p)$ is the associated relation for $(p, \underline{k}_1)$. $\square$

The correspondence on non-stable data (p, k_1) does not always preserve the associated relation in the same way as Lemma 4. Namely the correspondence is not involution. For example $(p, k_1) = (191, 15)$ is gotten the values $a = 22$, $n = 26$, $\tau = 0$, $q_2 = -73$. Then the image of $(191, 15)$ is $(102, 11)$, but the associated relation of $(102, 11)$ is $6k_1^2 - k_1 - 1 = 0(102)$ and $\tau = 1$. Moreover the image of $(102, 11)$ is $(87, 10)$ with the same associated relation as $(102, 11)$ and $\tau = 1$.

2.8 The difference of $A(x)$.

We define $A(x)$ to be

$$A(\epsilon_1\epsilon_2 a - 1, \epsilon_1 x, 1).$$

By using Formula (7), we calculate the difference of $A(x)$ in this section.

2.8.1 The $\epsilon_2 = 1$ case

Lemma 5 *Let p, k_1, a be the same parameters as above. Then the differences*

$$\begin{aligned} & A(\epsilon_1 a - 1, \epsilon_1 x, 1) - A(\epsilon_1 a - 1, \epsilon_1 x - 1, 1) \\ = \quad & E_{k_2}(\epsilon_1 x q_2 + k_2 + \epsilon_1 a) - E_{k_2}(\epsilon_1 x q_2 + \epsilon_1 a) \end{aligned} \tag{9}$$

are divided into 3 *cases as follows.*

(a) The condition (9)$= -1$ is equivalent to the following.
$(\epsilon_1 = 1)$ There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \tfrac{p\ell}{|q_2|} \rfloor + 1 \text{ or} \\ -q_1 j \ (p) \quad (j = 0, \cdots, a-1) \quad \left(\lfloor \tfrac{p\ell}{|q_2|} \rfloor \right). \end{cases}$$

$(\epsilon_1 = -1)$ There exists an integer ℓ such that

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \text{ and } \neq -q_1 j \ (p) \quad (j = 1, \cdots, a).$$

(b) The condition (9)$= 1$ is equivalent to the following.
$(\epsilon_1 = 1)$ There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \tfrac{p\ell}{|q_2|} \rfloor, \quad \text{and} \quad \neq -q_1 j \ (p) \quad (j = 0, 1, \cdots, a-1), \\ -q_1 j - 1 \ (p) \quad (j = 0, 1, \cdots, a-1), \quad (\lfloor \tfrac{p\ell}{|q_2|} \rfloor - 1), \text{ or} \\ -q_1 j + k_1 \ (p) \quad (j = 0, 1, \cdots, a-1), \quad (\lfloor \tfrac{p\ell}{|q_2|} \rfloor - 1). \end{cases}$$

$(\epsilon_1 = -1)$ There exists an integer ℓ such that

$$x = \begin{cases} -q_1 j \quad (p) \quad (j = 1, \cdots, a) \quad \left(\lfloor \tfrac{p\ell}{|q_2|} \rfloor + 1 \right), \text{ or} \\ \lfloor \tfrac{p\ell}{|q_2|} \rfloor \text{ and } \neq -q_1 j - 1 \ (p), \ -q_1 j - k_1 \quad (j = 1, \cdots, a). \end{cases}$$

(c) (9)$= 0$ otherwise.

Proof. (a) The condition (9)$= -1$ is equivalent to $p\ell < \epsilon_1 x q_2 + \epsilon_1 a \leq p\ell + k_2$

$$\Leftrightarrow x - 1 \leq \frac{p\ell}{|q_2|} < x + \frac{a}{q_2}.$$

If $\epsilon_1 = 1$ then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1$$

or

$$x = -q_1 j \quad (p) \quad (j = 0, \cdots, a-1) \quad (x = \lfloor \frac{p\ell}{|q_2|} \rfloor).$$

If $\epsilon_1 = -1$ then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \text{ and } \neq -q_1 j \quad (j = 1, \cdots, a).$$

(b) The condition (9)$= 1$ is equivalent to $p\ell < \epsilon_1 x q_2 + k_2 + \epsilon_1 a \leq p\ell + k_2$

$$\Leftrightarrow x + \frac{a}{q_2} \leq \frac{p\ell}{|q_2|} < x + 1 + \frac{2a}{q_2}.$$

If $\epsilon_1 = 1$ then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq -q_1 j \ \ (p) \ \ (j = 0, \cdots a-1)$$

or

$$x = -q_1 j - 1 \ (p) \ (j = 0, 1, \cdots, 2a-1).$$

If $\epsilon_1 = -1$ then this condition is

$$x = -q_1 j \ (p) \ (j = 1, \cdots, a) \quad \left(x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \right)$$

or

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq -q_1 j - 1 \ (p) \ (j = 1, \cdots, 2a).$$

□

2.8.2 The $\epsilon_2 = -1$ case

Lemma 6 *Let p, k_1, and a be the same parameters as above. Then the difference*

$$A(-\epsilon_1 a - 1, \epsilon_1 x, 1) - A(-\epsilon_1 a - 1, \epsilon_1 x - 1, 1) \tag{10}$$

is divided into three cases as follows.

(a) The condition (10)$= -1$ is equivalent to the following.
($\epsilon_1 = 1$) There exists an integer ℓ such that

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \ \text{ and } \neq q_1 j \ (p) \ \ (j = 1, \cdots, a).$$

($\epsilon_1 = -1$) There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \frac{p\ell}{|q_2|} \rfloor + 1, \text{ or} \\ q_1 j \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor \right). \end{cases}$$

(b) The condition (10)$= 1$ is equivalent to the following.
($\epsilon_1 = 1$) There exists an integer ℓ such that

$$x = \begin{cases} q_1 j \ (p) \ \ (j = 1, \cdots, a) \ \ (\lfloor \frac{p\ell}{|q_2|} \rfloor + 1), \text{ or} \\ \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq q_1 j - 1, \ \ q_1 j - k_1 \ (p) \ \ (j = 1, 2, \cdots, a). \end{cases}$$

($\epsilon_1 = -1$) There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq q_1 j \ (p) \ \ (j = 0, \cdots, a-1), \\ q_1 j - 1 \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor - 1 \right), \text{ or} \\ q_1 j + k_1 \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor - 1 \right). \end{cases}$$

(c) (10)$= 0$ otherwise.

Proof. **(a)** The condition (9)$= -1$ is equivalent to $p\ell < \epsilon_1 x q_2 - \epsilon_1 a \le p\ell + k_2$

$$\Leftrightarrow x - 1 \le \frac{p\ell}{|q_2|} < x - \frac{a}{q_2}.$$

If $\epsilon_1 = 1$ then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \ \text{ and } \ne q_1 j \ \ (j = 1, \cdots, a)$$

If $\epsilon_1 = -1$, then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1$$

or

$$x = q_1 j \ \ (j = 0, \cdots, a-1) \ \ \left(x = \lfloor \frac{p\ell}{|q_2|} \rfloor \right).$$

(b) This condition (9)$= 1$ is equivalent to $p\ell < \epsilon_1 x q_2 + k_2 - \epsilon_1 a \le p\ell + k_2$

$$\Leftrightarrow x - \frac{a}{q_2} \le \frac{p\ell}{|q_2|} < x + 1 - \frac{2a}{q_2}.$$

If $\epsilon_1 = 1$, then this condition is

$$x = q_1 j(p) \ \ (j = 1, \cdots, a) \ \ (x = \lfloor \frac{p\ell}{q_2} \rfloor + 1)$$

or

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor \ \text{and} \ \ne q_1 j - 1 \ \ (j = 1, \cdots, 2a).$$

If $\epsilon = -1$, then this condition is

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor \ \text{and} \ \ne q_1 j(p) \ \ (j = 0, \cdots, a-1)$$

or

$$x = q_1 j(p) \ \ (j = 0, \cdots, 2a-1) \ \ \left(x = \lfloor \frac{p\ell}{|q_2|} \rfloor - 1 \right).$$

□

As a result we have the following.

Lemma 7 *Let p, k_1, a be the same parameters as above. Then the difference*

$$A(\epsilon_1\epsilon_2 a - 1, \epsilon_1 x, 1) - A(\epsilon_1\epsilon_2 a - 1, \epsilon_1 x - 1, 1) \qquad (11)$$

is divided into three cases as follows.

(a) The condition (11)$= -1$ is equivalent to the following.
($\epsilon_1\epsilon_2 = 1$) There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \frac{p\ell}{|q_2|} \rfloor + 1, \text{ or} \\ -\epsilon_2 q_1 j \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor \right). \end{cases}$$

($\epsilon_1\epsilon_2 = -1$) There exists an integer ℓ such that

$$x = \lfloor \frac{p\ell}{|q_2|} \rfloor + 1 \ \text{ and } \ne -\epsilon_2 q_1 j \ (p) \ \ (j = 1, \cdots, a).$$

(b) The condition (11)$= 1$ is equivalent to the following.
($\epsilon_1\epsilon_2 = 1$) There exists an integer ℓ such that

$$x = \begin{cases} \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq -\epsilon_2 q_1 j \ (p) \ \ (j = 0, \cdots, a-1), \\ -\epsilon_2 q_1 j - 1 \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor - 1 \right), \text{ or} \\ -\epsilon_2 q_1 j + k_1 \ (p) \ \ (j = 0, \cdots, a-1) \ \ \left(\lfloor \frac{p\ell}{|q_2|} \rfloor - 1 \right). \end{cases}$$

($\epsilon_1\epsilon_2 = -1$) There exists an integer ℓ such that

$$x = \begin{cases} -\epsilon_2 q_1 j \ (p) \ \ (j = 1, \cdots, a) \ \ (\lfloor \frac{p\ell}{|q_2|} \rfloor + 1), \text{ or} \\ \lfloor \frac{p\ell}{|q_2|} \rfloor \text{ and } \neq -\epsilon_2 q_1 j - 1, \ \ -\epsilon_2 q_1 j - k_1 \ (p) \ \ (j = 1, 2, \cdots, a). \end{cases}$$

(c) (11)$= 0$ otherwise.

3 The global view of the function $A(i + jk_1)$

3.1 The local behavior of $A(x)$

From Lemma 7, x in which $A(x) \neq A(x - \epsilon_1)$ is $x = \lfloor \frac{p\ell}{|q_2|} \rfloor - 1, \lfloor \frac{p\ell}{|q_2|} \rfloor, \lfloor \frac{p\ell}{|q_2|} \rfloor + 1$. We consider the quotient

$$\frac{p}{|q_2|} = \tau + \frac{\gamma' + \epsilon_1\epsilon_2}{n} + \frac{a}{n|q_2|}. \tag{12}$$

Here we put $\alpha = \gamma' + \epsilon_1\epsilon_2 + \frac{a}{|q_2|}$. In particular we have

$$\lfloor \frac{np}{|q_2|} \rfloor = n\tau + \gamma' + \epsilon_1\epsilon_2 + \lfloor \frac{a}{|q_2|} \rfloor = k_1 + \epsilon_1\epsilon_2.$$

3.1.1 The $\epsilon_1\epsilon_2 = 1$ case.

Since by Theorem 3 the values of $\tilde{a}_i$ are 0, ±1, or 2, the possibilities of the local behavior of $A(x)$ around $x = \lfloor \frac{p\ell}{|q_2|} \rfloor$ are Figure 1, 2, and 3 in the case of $\epsilon_1 = 1$, and Figure 6, 7, and 8 in the case of $\epsilon_1 = -1$.

Now suppose that $\epsilon_1 = 1$. If $A(x_0) = 2$ holds for some x_0, then we cannot find any function $A(x)$ by connecting Figure 1, 2, or 3. For example if the type (+,iii) appears, then the first possible type which is non-constant with $x > x_0$ is (+,v) or (+,iii). Since the integer x with $A(x) = 2$ is unique in $\mathbb{Z}/p\mathbb{Z}$, $\Delta_K(t) = x^{-1} + 1 + x$ holds. This polynomial cannot be any Alexander polynomial of a knot in a homology sphere. Hence we may assume that $|A(x)| \leq 1$ holds and (+,i), (+,ii), (+,iv), or (+,vi) are applicable non-constant local behaviors of $A(x)$.

We put $\alpha(j) = -\epsilon_2 q_1 j$ and $\beta(j) = -\epsilon_2 q_1 j + k_1$. In the case of $\epsilon_1 = 1$, we can get Figure 4, in which there are the values of $A(x)$ from $\alpha(j) - 1$ to $\beta(j) + 1$, fitting together of local behaviors for the values to be connected.



$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor - 1$	$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor$	$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor + 1$	x
0	1	0	(+,i)
-1	0	-1	(+,ii)
1	2	1	(+,iii)

Figure 1: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, 1)$, $\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor \neq -\epsilon_2 q_1 j$



$-\epsilon_2 q_1 j - 2$	$-\epsilon_2 q_1 j - 1$	$-\epsilon_2 q_1 j$	$-\epsilon_2 q_1 j + 1$	x
0	1	0	-1	(+,iv)
1	2	1	0	(+,v)

Figure 2: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, 1), j = 0, \cdots, a - 1)$



$-\epsilon_2 q_1 j$ $+k_1 - 1$	$-\epsilon_2 q_1 j$ $+k_1$	$-\epsilon_2 q_1 j$ $+k_1 + 1$	$-\epsilon_2 q_1 j$ $+k_1 + 2$	x
-1	0	1	0	(+,vi)
0	1	2	1	(+,vii)

Figure 3: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, 1), j = 0, \cdots, a - 1)$

x	$\alpha(j)-1$	$\alpha(j)$	$\alpha(j)+1$	$\cdots$	$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor-1$	$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor$	$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor+1$	$\cdots$	$\cdots$	$\lfloor\frac{p(\ell+1)}{\lvert q_2\rvert}\rfloor-1$	$\lfloor\frac{p(\ell+1)}{\lvert q_2\rvert}\rfloor$	$\lfloor\frac{p(\ell+1)}{\lvert q_2\rvert}\rfloor+1$	$\cdots$	$\beta(j)-1$	$\beta(j)$	$\beta(j)+1$
$A(x)$	1	0	-1	$\cdots$	-1	0	-1	$\cdots$	$\cdots$	-1	0	-1	$\cdots$	-1	0	1

Figure 4: The values $A(x)$ from $\alpha(j)-1$ to $\beta(j)+1$ in the case of $(\epsilon_1,\epsilon_2)=(1,1)$.

In the region $\alpha(j)+1\le x\le\beta(j)-1$ there are 0 and -1. We call the integer points in the region *negative region.*

On the other hand Figure 5 is the values of $A(x)$ between $\beta(j)-1$ and $\alpha(k)+1$.

x	$\beta(j)-1$	$\beta(j)$	$\beta(j)+1$	$\beta(j)+2$	$\cdots$	$\lfloor\frac{p\ell'}{\lvert q_2\rvert}\rfloor-1$	$\lfloor\frac{p\ell'}{\lvert q_2\rvert}\rfloor$	$\lfloor\frac{p\ell'}{\lvert q_2\rvert}\rfloor+1$	$\cdots$	$\cdots$	$\lfloor\frac{p(\ell'+1)}{\lvert q_2\rvert}\rfloor-1$	$\lfloor\frac{p(\ell'+1)}{\lvert q_2\rvert}\rfloor$	$\lfloor\frac{p(\ell'+1)}{\lvert q_2\rvert}\rfloor+1$	$\cdots$	$\alpha(k)-2$	$\alpha(k)-1$	$\alpha(k)$	$\alpha(k)+1$
$A(x)$	-1	0	1	0	$\cdots$	0	1	0	$\cdots$	$\cdots$	0	1	0	$\cdots$	0	1	0	-1

Figure 5: The values $A(x)$ from $\beta(j)-1$ to $\alpha(j)+1$ in the case of $(\epsilon_1,\epsilon_2)=(1,1)$.

Then the values $A(x)$ between $\beta(j)+1$ and $\alpha(k)-1$ are 0 or 1. We call the integer points in this region *positive region.*

Thus if (p,k_1) is a data realized by a lens space surgery over S^3 or $\Sigma(2,3,5)$, then we can decide the values $A(x)$ completely. Hence we figure out that each positive region and negative region appear alternatively and do not overlap each other.

In the same way all the local behaviors with $\epsilon_1=-1$ are from Figure 6, 7, and 8. The case satisfying $A(x)=2$ for some integer x is classified in Section 3.1.3.

Thus we can decide the values $A(x)$ for any x. Thus we call the integer points between $\alpha(j)+2$ and $\beta(j)$ and integer points between $\beta(j)+3$ and $\alpha(k)$ *positive region* and *negative region* respectively.

Next we put the values $A(i+jk_1)$ on i-j plain. We call this distribution *global view* of the Alexander polynomial. Since we have

$$\lfloor\frac{mnp}{|q_2|}\rfloor=mn\tau+m(\gamma'+\epsilon_1\epsilon_2)+\lfloor\frac{ma}{|q_2|}\rfloor=m(k_1+\epsilon_1\epsilon_2)+\lfloor\frac{ma}{|q_2|}\rfloor.$$

Then the following Lemma is true.

Lemma 8

$$\begin{aligned}\lfloor\frac{mnp}{|q_2|}\rfloor-\lfloor\frac{(m-1)np}{|q_2|}\rfloor-(k_1+\epsilon_1\epsilon_2) &= \lfloor\frac{ma}{|q_2|}\rfloor-\lfloor\frac{(m-1)a}{|q_2|}\rfloor \\ &= \begin{cases}1 & 0\le[ma]_{|q_2|}<a\\ 0 & \textit{otherwise}\end{cases}\end{aligned}\tag{13}$$

Since we have $a'=a^{-1}=-\epsilon_1(k_1-\epsilon_1\epsilon_2)\ \ (|q_2|)$, when $m=a'i\ \ (i=0,\cdots,a-1)$, we have (13) $=1$. Hence we have

$$\lfloor\frac{a'inp}{|q_2|}\rfloor=-\epsilon_1 iq_1\ \ (p).$$



$\lfloor \frac{p\ell}{\lvert q_2\rvert} \rfloor$	$\lfloor \frac{p\ell}{\lvert q_2\rvert} \rfloor + 1$	$\lfloor \frac{p\ell}{\lvert q_2\rvert} \rfloor + 2$	x
1	0	1	(+,iix)
0	-1	0	(+,ix)

Figure 6: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, -1)$, $\lfloor \frac{p\ell}{|q_2|} \rfloor \neq -\epsilon_2 q_1 j$



$-\epsilon_2 q_1 j - 1$	$-\epsilon_2 q_1 j$	$-\epsilon_2 q_1 j + 1$	$-\epsilon_2 q_1 j + 2$	x
0	-1	0	1	(+,x)
1	0	1	2	(+,xi)

Figure 7: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, -1)$, $j = 0, \cdots, a-1$)



$-\epsilon_2 q_1 j$ $+k_1$	$-\epsilon_2 q_1 j$ $+k_1 + 1$	$-\epsilon_2 q_1 j$ $+k_1 + 2$	$-\epsilon_2 q_1 j$ $+k_1 + 3$	x
1	0	-1	0	(+,xii)
2	1	0	1	(+,xiii)

Figure 8: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, -1)$,($j = 0, \cdots, a-1$)

This implies that if $\epsilon_1 = 1$ (or $\epsilon_1 = -1$), then all x that gives $A(x) = 1$ (or $A(x) = -1$) lines like stairs in positive region (or negative region) on i-j plain. See Figure 9. Here each of places which put $+$ or $-$ represents $A(x) = 1$ or -1 respectively and the blanks are all 0. The starting point of a stair-like sequence is (x, y), where $(i, j) = (x-1, y-1)$ is a point in negative (or positive) region that $A(i + jk_1)$ is 0, and the end point is (z, w), where $(i, j) = (z + 1, w + 1)$ is a point in negative (or positive) region that $A(i + jk_1)$ is 0.



$\cdots - \ 0 \ - \cdots - \ - \ - \ - \ 0 \ - \cdots$
positive region
$+$
$\cdots - \ 0 \ - \ - \cdots \quad \cdots - \ 0 \ - \ - \cdots\cdots - \ 0 \ - \cdots$

Figure 9: The stair-like sequences in positive region. ($\epsilon_1 = 1$)

Combining the results above, we plot $A(i + jk_1)$ on i-j-plane to get Figure 10, 11 in the case of $\epsilon_1 = 1$, -1 respectively.



j
$\beta(*)$
$\alpha(*)$
negative region
positive region
$0 \ 0 \ 0 \ + \ 0 \cdots$
$- \ 0 \ + \ 0 \cdots$
$+ \ 0 \ - \ - \cdots \cdots - \ 0 \ - \cdots - \ 0$
0
1
2
3
-1
0
1
2
$\lfloor \frac{p}{|q_2|} \rfloor$
$\lfloor \frac{2p}{|q_2|} \rfloor$
i

Figure 10: The global view of $A(i + jk_1)$ in the case of $(\epsilon_1, \epsilon_2) = (1, 1)$

3.1.2 The $\epsilon_1\epsilon_2 = -1$ case.

In this section we give the global view of $A(i+jk_1)$ by investigating the local behavior of $A(x)$ in the case of $\epsilon_1\epsilon_2 = -1$.

In the case of $\epsilon_1 = 1$, using Lemma 7, we can easily show that the possible non-constant local behaviors of the function $A(x)$ are Figure 12, 13, 14. In the case of $\epsilon_1 = -1$ the non-constant local behaviors are Figure 19, 20, 21. Here $\gamma(j)$ stands for $-\epsilon_2 q_1 j - k_1$. The case where $A(\alpha(j)) = 2$ for some integer j will be dealt with in Section 3.1.4. We assume that $|A(x)| = 1$ holds for any x this case will be dealt in Section 3.1.5.


j
$\beta(*)+1$
$+ \cdots + \; 0 \; - \; 0 \; \cdots$
$\cdots \; 0 \; - \; 0 \; + \; + \; \cdots$
positive region
$\alpha(*)+1$
$\beta(*)+1$
$+ \; + \; + \; + \; 0 \; \cdots \; + \; 0 \; -$
$\cdots \; 0 \; - \; 0 \; \cdots \; \cdots \; 0 \; - \; 0 \; +$
$\lfloor \frac{\ell p}{|q_2|} \rfloor + 1$
$\alpha(*)+1$
negative region
3
2
1
0
$0 \; 0 \; 0 \; - \; 0 \; \cdots$
$+ \; 0 \; - \; 0 \; \cdots$
$- \; 0 \; + \; + \; \cdots \; \cdots \; + \; 0 \; + \; \cdots \; + \; 0$
0
1
2
3
$\lfloor \frac{p}{|q_2|} \rfloor + 1$
$\lfloor \frac{2p}{|q_2|} \rfloor + 1$
i

Figure 11: The global view of $A(i + jk_1)$ in the case of $(\epsilon_1, \epsilon_2) = (-1, -1)$

$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor - 1$	$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor$	$\lfloor \frac{p\ell}{\lvert q_2 \rvert} \rfloor + 1$	x
0	1	0	(-,i)
-1	0	-1	(-,ii)
1	2	1	(-,iii)

$+1$ -1

Figure 12: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, -1)$, $\lfloor \frac{p\ell}{|q_2|} \rfloor \neq -\epsilon_2 q_1 j$ $(0 \leq j < a)$

$-\epsilon_2 q_1 j - 2$	$-\epsilon_2 q_1 j - 1$	$-\epsilon_2 q_1 j$	x
-1	-1	0	(-,iv)
0	0	1	(-,v)
1	1	2	(-,vi)

0 $+1$

Figure 13: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, -1)$, $(j = 1, \cdots, a)$

$-\epsilon_2 q_1 j$ $-k_1 - 1$	$-\epsilon_2 q_1 j$ $-k_1$	$-\epsilon_2 q_1 j$ $-k_1 + 1$	x
1	1	0	(-,vii)
0	0	-1	(-,iix)

0 -1

Figure 14: $A(x)$: $(\epsilon_1, \epsilon_2) = (1, -1)$, $(j = 1, \cdots, a)$

3.1.3 The great common divisor $\gcd(a, n)$.

We put $d = (\eta, a) = (n, a)$, $\eta = \eta_1 d$, $a = a_1 d$. We put $\tilde{\eta} := [\eta_1^{-1}]_{a_1}$. Let j be $j_1 a_1 + [-\epsilon_2 \tilde{\eta} j_2]_{a_1}$ with $0 \leq j_1 < d$, $0 \leq j_2 < a_1$. Then we have $-\epsilon_2 j q_1 = -\epsilon_2 \frac{j\eta p - \epsilon_1 j k_1 - \epsilon_2 j}{a} \equiv \frac{j_2}{a_1} p + \frac{j(\epsilon_1 \epsilon_2 k_1 + 1)}{a}$ (p). We define intervals to be

$$I_{j_1, j_2} := [-\epsilon_2(\frac{j_2}{a_1} p + \frac{j(\epsilon_1 \epsilon_2 k_1 + 1)}{a}), -\epsilon_2(\frac{j_2}{a_1} p + \frac{j(\epsilon_1 \epsilon_2 k_1 + 1)}{a} + k_1)]$$

if $\epsilon_1 \epsilon_2 = 1$, and

$$I_{j_1, j_2} := [-\epsilon_2(\frac{j_2}{a_1} p + \frac{j(\epsilon_1 \epsilon_2 k_1 + 1)}{a} - k_1), -\epsilon_2(\frac{j_2}{a_1} p + \frac{j(\epsilon_1 \epsilon_2 k_1 + 1)}{a})]$$

if $\epsilon_1 \epsilon_2 = -1$.

For a fixed j_2 the intervals I_{j_1, j_2} $(j_1 = 0, \cdots, d-1)$ are overlapped each other. Furthermore $I_{j_1, j_2} \cap I_{j_1, j_3} = \emptyset$ holds unless $j_2 = j_3$. On the other hand if $d \geq 2$, then we have $I_{j_1, j_2} \cap I_{j_3, j_4} = \emptyset$ unless $j_1 = j_3$, since $\frac{p}{a_1} \geq \frac{2p}{a} \geq \frac{2(ak_1^2 + \epsilon_1 k_1 + \epsilon_2)}{an} \geq 2k_1(\tau + \frac{\gamma'}{n}) - \frac{2(k_1+1)}{an} > 4k_1 - (k_1 + 1) = 3k_1 - 1$.

Thus $A(x)$ increases by $-\epsilon_2$ around $x = -\epsilon_2 j q_1$ as shown in Figure 2, 7, 13, and 20. In the same way $A(x)$ increases by ϵ_2 around $x = -\epsilon_2 j q_1 + \epsilon_1 \epsilon_2 k_1$ as it is shown in Figure 3, 8, 14, and 21. Therefore $d \leq 2$ holds by Theorem 3. Thus we showed the following.

Lemma 9 *Let* (p, k_1) *be a data realized by lens space surgery* S^3 *or* $\Sigma(2,3,5)$. *Then the parameters* a, n *have* $\gcd(a, n) = 1$ *or* 2.

3.1.4 The $2g(p, k_1) = p$ case

For the case where $2g = p$ is satisfied the case of $(\epsilon_1, \epsilon_2) = (-1, -1)$ and $2g = p$ is remained. If $A(x_0) = 2$ for some integer x, then using Figure 7 and 8 we can get the following for some integer j.

$-\epsilon_2 q_1 j - 1$	$-\epsilon_2 q_1 j$	$-\epsilon_2 q_1 j + 1$	$-\epsilon_2 q_1 j + 2$	$-\epsilon_2 q_1 j + 3$	$-\epsilon_2 q_1 j + 4$	$-\epsilon_2 q_1 j + 5$
1	0	1	2	1	0	1

Namely $-\epsilon_2 q_1 j + 2 = -\epsilon_2 q_1 j' + k_1$ holds.

In this case we have $d = 2$ and the local view around (i, j) satisfying $x = i + jk_1$ and $A(x) = 2$ is Figure 15. Since this pattern does not appear in other places up to the moving by $p\mathbb{Z}$, $a_1 = 1$ holds. Thus $I_{0,0} \cap I_{1,0}$ is overlapped as in Figure 16. Thus



Figure 15: The local view of x with $A(x) = 2$.



Figure 16: The overlapped interval.

we have $-\epsilon_2 q_1 j + 2 = q_1 + 2 = k_1$. From the associated relation $2k_1^2 - k_1 - 1$, we get $k_1 = 5$ and $q_1 = 3$. Thus we have $p|22$. If $p = 11$, we cannot find lens space surgery from the list in [8]. Therefore we have $p = 22$. This case is realized by lens space surgery of type E_2 and $J = -1$ in Table 2

In the case $\epsilon_1\epsilon_2 = -1$ we can show in the same way as above that the $2g = p$ case does not exist.

3.1.5 The global view in the case of $\epsilon_1\epsilon_2 = -1$

We get back to the description of the global view. Here we suppose that $d = 2$ holds. In particular a is even number. Then we get the values as in Figure 17. Note that if $\lfloor \frac{p\ell}{|q_2|} \rfloor$ appears between $\alpha(j_2)$ and $\gamma(a_1 + j_2 + 1)$, it is at most one time. Then we can reduce the following inequality.

x	$\gamma(a_1+j_2)$	$\gamma(a_1+j_2)+1$	$\cdots$	$\gamma(j_2)-1$	$\gamma(j_2)$	$\gamma(j_2)+1$	$\cdots$	$\alpha(a_1+j_2)-2$
$A(x)$	1	0	$\cdots$	0	0	-1	$\cdots$	-1

$\alpha(a_1+j_2)-1$	$\alpha(a_1+j_2)$	$\cdots$	$\alpha(j_2)-2$	$\alpha(j_2)-1$	$\alpha(j_2)$	$\cdots$	$\gamma(a_1+j_2+1)$
-1	0	$\cdots$	0	0	1	$\cdots$	1

Figure 17: The values $A(x)$ from $\gamma(a_1 + j_2)$ to $\gamma(a_1 + j_2 + 1)$ in the case of $d = 2$ and $\epsilon_1\epsilon_2 = -1$



Figure 18: The $d = 2$ case.

$$ak_1 + (\tau + 2)\frac{a}{2} + \tau > p = \frac{ak_1^2 + \epsilon_1 k_1 + \epsilon_2}{n}$$

Thus

$$ak_1(\tau - 1 + \frac{\gamma'}{n}) < \frac{k_1 + 1}{n} + \frac{n\tau + \gamma'}{n}\frac{a}{2} + a + \tau < \frac{k_1}{n}(2 + \frac{a}{2}) + \frac{1}{n} + a$$

From $\frac{2}{an} \leq \frac{1}{n}$ and $\frac{1}{k_1} < \frac{1}{2n}$,

$$< ak_1(\frac{2}{an} + \frac{1}{2n} + \frac{1}{k_1}) + \frac{1}{n} < ak_1\frac{2}{n} + \frac{1}{n}$$

Therefore

$$ak_1(\tau - 1 + \frac{\gamma' - 2}{n}) < \frac{1}{n}$$

Since the left hand side is more than one, this inequality is inconsistent.

Therefore we must have $d = 1$, namely $(a, n) = 1$. If $A(\gamma(j)) = 1$ holds for some j, then $A(x) \leq 0$ holds for all x. This is equivalent to $\Delta_{p,k_1}(t) = 1$ and the data is realized by the unknot surgery. Therefore the non-constant local possibilities are (-,i), (-,ii), (-,iv) and (-,iix).

In the case of $\epsilon_1 = -1$, we can deduce that the non-constant local behaviors are (-,ix), (-,x), (-,xi), (-,xiv) in the same argument as in the $\epsilon_1 = 1$ case.



$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor$	$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor + 1$	$\lfloor\frac{p\ell}{\lvert q_2\rvert}\rfloor + 2$	x
1	0	1	(-,ix)
0	-1	0	(-,x)

Figure 19: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, 1)$, $\lfloor\frac{p\ell}{|q_2|}\rfloor \neq -\epsilon_2 q_1 j$ $(0 \leq j < a)$



$-\epsilon_2 q_1 j - 1$	$-\epsilon_2 q_1 j$	$-\epsilon_2 q_1 j + 1$	x
1	1	0	(-,xi)
0	0	-1	(-,xii)

Figure 20: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, 1)$, $(j = 1, \cdots, a)$



$-\epsilon_2 q_1 j$ $-k_1$	$-\epsilon_2 q_1 j$ $-k_1 + 1$	$-\epsilon_2 q_1 j$ $-k_1 + 2$	x
-1	-1	0	(-,xiii)
0	0	1	(-,xiv)

Figure 21: $A(x)$: $(\epsilon_1, \epsilon_2) = (-1, 1)$, $(j = 1, \cdots, a)$

Combining these results, we plot $A(i + jk_1)$ on i-j-plane to get Figure 22, 23 in the case of $\epsilon_1 = 1$, -1 respectively.

4 A certain integrality condition

Here for any $1 \leq i \leq a - 1$ we define n_i to be $[-\epsilon_2 \eta^{-1} i]_a$, so that we have

$$-\epsilon_2 n_i q_1 = \frac{ip + n_i(\epsilon_1 \epsilon_2 k_1 + 1)}{a} \quad (p). \tag{14}$$



α(∗)
j
− ⋯ − 0 0 0 ⋯
⋯ 0 0 0 − − ⋯
negative region
γ(∗)
α(∗)
0 − − − − − 0 ⋯ − 0 0
+
0 0 −
0 +
positive region
1 + 0 +
γ(∗)
0 0 + 0 +
−1 − − 0 ⋯ 0 + 0 +
−2 0 − − ⋯ − 0 − ⋯ − 0
−1 0 1 2

Figure 22: The global view of $A(i + jk_1)$ in the case of $(\epsilon_1, \epsilon_2) = (1, -1)$



α(∗)
j
+ ⋯ + 0 0 0 ⋯
⋯ 0 0 0 + + ⋯
positive region
γ(∗)
α(∗)
+ 0 + ⋯ + 0 0 ⋯
− ⋯ 0 0 + ⋯
negative region
−
γ(∗)
−
−1 + + 0 0 ⋯ ⋯ 0 − 0 ⋯
−2 0 0 + ⋯ + + 0
0 1 2
⌊p/|q₂|⌋ + 1
i

Figure 23: The global view of $A(i + jk_1)$ in the case of $(\epsilon_1, \epsilon_2) = (-1, 1)$

In the case of $\epsilon_1\epsilon_2 = 1$ we put $n_0 = 0$ then $-\epsilon_2 n_0 q_1 = 0$. In the case of $\epsilon_1\epsilon_2 = -1$ we put $n_a = a$ then $-\epsilon_2 n_a q_1 = \epsilon_1\epsilon_2 k_1 + 1 = -k_1 + 1$.

We define n' to be the integer satisfying $an' = nn_{a-1} - \epsilon_2$. Then $a(n-n') = nn_1 + \epsilon_2$ holds. We define n'_i to be the integer satisfying $an'_i = nn_i + \epsilon_2 i$ $(1 \leq i \leq a-1)$. When $\epsilon_1\epsilon_2 = 1$, we have $n'_0 = 0$. When $\epsilon_1\epsilon_2 = -1$, we have $n'_a = n + \epsilon_2$. Then $n'_1 = n - n'$ and $n'_{a-1} = n' + \epsilon_2$ holds. We shall show the integral condition.

Lemma 10 (integrality condition) *Let γ', n_1 be as above. Then (14) is integer for any $i = 1, 2, \cdots, a-1$ if and only if*

$$(\gamma')^2 - \epsilon_1\epsilon_2 n'\gamma' - n' = 0 \quad (n)$$

holds.

Proof. From the integrality of (14) we have

$$\begin{aligned}(14) &= in\tau^2 + 2i\gamma'\tau + \frac{i(\gamma')^2 + n'_i(\epsilon_2 k_1 + 1)}{n} \\ &= in\tau^2 + (2i\gamma' + \epsilon_1\epsilon_2 n'_i)\tau + \frac{i(\gamma')^2 + \epsilon_1\epsilon_2 n'_i\gamma' + n'_i}{n}.\end{aligned}$$

Thus from $n'_i = in'_1$ (n) for any i we have

$$i(\gamma')^2 + \epsilon_1\epsilon_2 n'_i\gamma' + n'_i = i((\gamma')^2 + \epsilon_1\epsilon_2 n'_1\gamma' + n'_1) = 0 \quad (n).$$

Therefore the integrality of (14) is equivalent to

$$(\gamma')^2 + \epsilon_1\epsilon_2 n'_1\gamma' + n'_1 = 0 \quad (n).$$

□

5 Classification

In this section we classify all stable data realized lens space surgery on S^3 or $\Sigma(2,3,5)$ under the conditions $a \geq 2$, $0 < \gamma' < n$, $k_1 \geq 4$, $n \geq 2$ and $g(p, k_1) < \frac{p}{2}$.

Proposition 2 *If the pair (p, k_1) is realized by a surgery over S^3 or $\Sigma(2,3,5)$ and $n = 1$ holds, then $(k_1, k_2) = 1$ or 2 and the surgery is realized by (I), (II).*

Proof From Theorem 12 in [7] $(k_1, k_2) = 1$ or 2 holds. If $(k_1, k_2) = 1$, then (p, k_1) is realized by (I) and if $(k_1, k_2) = 2$, then (p, k_1) is realized by (II). □

5.1 The case of $\gamma' = n - 1$ or 1.

Hence for any integer m we have

$$\lfloor \frac{mp}{|q_2|} \rfloor - \lfloor \frac{(m-1)p}{|q_2|} \rfloor = \tau + \lfloor \frac{\alpha m}{n} \rfloor - \lfloor \frac{\alpha(m-1)}{n} \rfloor. \quad (15)$$

Since $0 < \frac{\alpha}{n} < 2$, (15) is τ, $\tau + 1$ or $\tau + 2$. If for some integer m the value of (15) is $\tau + 2$, then $\alpha > n$, $\gamma' \geq n - 1$, and $\epsilon_1\epsilon_2 = 1$ hold.

Lemma 11 *If (15) is τ+2, then the stable data (p, k_1) is realized by lens space surgery type (I) or (II).*

Proof. If (15) is $\tau+2$, we have $\alpha=\gamma'+\epsilon_1\epsilon_2+\frac{a}{|q_2|}>n$. From $\gamma'<n$, we have $\gamma'=n-1$ and $\epsilon_1\epsilon_2=1$. From integral condition $1-n'+n'=1=0(n)$, $n=1$ holds. This data (p,k_1) is realized by Dehn surgeries of (I) or (II) by Proposition 2. □

If $\gamma'=n-1$ and $\epsilon_1\epsilon_2=-1$, then from the integrality condition $1-n'-n'=1-2n'=0(n)$ we have $a=-2\epsilon_2=2\epsilon_1\ (n)$. Thus we can put $a=un+2\epsilon_1$ for some integer u. Therefore $k_1=n\tau+n-1$ and

$$\begin{aligned} p &= (un+2\epsilon_1)n\tau^2+(2a(n-1)+\epsilon_1)\tau+an-2na+\epsilon_1+u \\ &= u(n\tau+n-1)^2+\epsilon_1(2n\tau^2+(4n-3)\tau+2n-3) \\ &= uk_1^2+\epsilon_1(2k_1-1)(\tau+1) \end{aligned}$$

Since $(n,a)=1$ holds, n is odd number. Each of the data is realized by type (III_+) by Berge's list. Actually put $J=\epsilon_1(\tau+1)$.

In the case of $\gamma'=1$ in the same way, if we have $\epsilon_1\epsilon_2=1$, we can get

$$(p,k_1)=(uk_1^2-\epsilon_1(2k_1+1)\tau,n\tau+1)$$

for some integer u, where n is odd. If $\epsilon_1\epsilon_2=-1$ then $n=1$.

The data of the former part is realized by type (III_-). (Put $J=-\epsilon_1\tau$. The data of the latter part is realized by type (I) or (II).

5.2 Forbidden and admitted sequences

Definition 4 *Let $m_i (i\in\mathbb{Z})$ be an integer-valued sequence. We call a subsequence $m_{i_0},m_{i_0+1},\cdots,m_{i_1}$ satisfying $m_{i_0}=m_{i_1}=-1$ and $m_{i_0+j}=0$ $(1\le j\le i_1-i_0-1)$ forbidden subsequence. We call a subsequence $m_{i_0},m_{i_0+1},\cdots,m_{i_1}$ satisfying $m_{i_0}=m_{i_1}=1$ and $m_{i_0+j}=0$ $(1\le j\le i_1-i_0-1)$ admitted subsequence.*

If the sequence $\{\tilde{a}_i\}$ includes a forbidden subsequence, this data is not realized by any lens space surgery over S^3 or $\Sigma(2,3,5)$. If the sequence $\{\tilde{a}_i\}$ includes an admitted subsequence and the data is realized by a lens space surgery over S^3 or $\Sigma(2,3,5)$, then this subsequence $\tilde{a}_{i_0},\cdots,\tilde{a}_{i_1}$ satisfies $i_0=g(p,k_1)\ (p)$, $i_1=p-g(p,k_1)\ (p)$ and $i_1-i_0=p-2g(p,k_1)$.

5.3 Blocks

We consider polygons surrounded by a bold dotted line in the global view on i-j-plane as in Figure 24. This polygon is composed of 6 segments with lattice point $\mathbb{Z}^2$ as the ends of the segments. The top and bottom segments are the maximal one that all the lattice points x on it have $A(x)=\pm1$. Let $\{(x,y)|y=s_0,t_0\le x\le t_1\}$ and $\{(x,y)|y=s_1,b_0\le x\le b_1\}$ be the top and bottom line respectively. The two of the remaining 4 lines are

$$\{(x,y)|y=t_1,\ s_1-1\le y\le s_1\},\ \{(x,y)|x=b_0,\ s_0\le y\le s_0+1\},$$

or

$$\{(x,y)|x=t_0,s_1-1\le y\le s_1\},\ \{(x,y)|x=b_1,\ s_0\le y\le s_0+1\}.$$

for $\epsilon_1\epsilon_2=1$ or -1 respectively. These make two components. The more two segments are two lines constructed by connecting each of right end points of the components and each of left end points of that. Furthermore t_0,t_1,b_0,b_1,s_0, and s_1 satisfy

$$t_1-b_0=s_1-s_0-2$$

or

$$b_1 - t_0 = s_1 - s_0 - 2.$$

The former appears in global view of $\epsilon_1\epsilon_2 = 1$ and the latter in global view of $\epsilon_1\epsilon_2 = -1$. Note that here are some ±1 on line connecting (t_1, s_0-1) and (b_0, s_1) or (t_0, s_0-1) and (b_1, s_1+1), where each ±1 on the line has opposite sign on top and bottom line. We call this polygon *a block*. Let denote by $B_{i,j}$ a block with the (i,j) component as in Figure 25. However actually in the case of $\epsilon_1\epsilon_2 = -1$ the bottom line of $B_{0,0}$ is not lies on the i-axis. One must note that the accurate location of blocks in the case later. The left one in Figure 24 is called (+)-block and the right one (-)-block. Here there are 1 or -1 in the top or bottom row of the block. The points except the top, bottom and left (or right) sides of (+)-block (or (-)-block) have all 0.



Figure 24: Blocks



Figure 25: $B_{i,j}$:(i,j) component of blocks

We define *top width* $\partial_+B_{i,j}$ for a block to be $t_1 - t_0 + 1$ and *bottom width* $\partial_-B_{i,j}$ to be $b_1 - b_0 - 1$, and define *height* $H(B_{i,j})$ to be $s_1 - s_0 + 1$. The right hand side of (15) calculates the top (or bottom) width of the block.

There exists a point (i_1, j_1) satisfying $i_1 + j_1k = \lfloor \frac{p\ell}{|q_2|} \rfloor$ around the vertex of the right (or left) top (or bottom) in a block $B_{i,j}$. We denote the points by $\partial^2_{lt}B_{i,j}$, $\partial^2_{rt}B_{i,j}$, $\partial^2_{lb}B_{i,j}$, and $\partial^2_{rb}B_{i,j}$ corresponding to the left top, right top, left bottom, and right bottom vertices respectively as in Figure 27, in which the center points of the circles are the points $\partial^2_{*}B_{i,j}$ for $* = lt, rt, lb$, and rb.

Figure 26 presents examples of the blocks with the height $\tau - 1$, τ, or $\tau + 1$. top width and bottom width is τ or $\tau + 1$.

As a result the next lemma easily follows.


$H=\tau-1$
$H=\tau$
$H=\tau+1$

Figure 26: blocks


$\partial_t B_{i,j}$
$\partial^2_{lt} B_{i,j}$
$\partial^2_{rt} B_{i,j}$
$B_{i,j}$
$\partial^2_{lb} B_{i,j}$
$\partial^2_{rb} B_{i,j}$
$\partial_b B_{i,j}$

Figure 27: The vertices of a block $B_{i,j}$. $(\epsilon_1, \epsilon_2) = (-1, 1)$

Lemma 12 *The top (or bottom) width of any block is τ or $\tau+1$. Furthermore if top (or bottom) width of $B_{i_0,j}$ and $B_{i_0+m,j}$ are $\tau+1$ and the top (or bottom) width of $B_{i_0+i_1,j}$ for any $1 \le i_1 \le m-1$ is τ, then $m = \lfloor \frac{n}{\alpha} \rfloor$ or $1 + \lfloor \frac{n}{\alpha} \rfloor$. These points are called the 2nd boundary of the block. In particular, the top width of $B_{-1,0}$ is $\tau+1$*

Proof. We show just the last assertion. Since $[\alpha 0]_n = 0$ holds, Lemma 13 implies $B_{-1,0}$ has $\tau+1$ top width. $\square$

5.4 The $2 \le \gamma' \le n-2$ case (i).

From this section we assume that $2 \le \gamma' \le n-2$ holds. In particular $n \ge 4$, $\alpha < n$, and

$$\lfloor \frac{p}{|q_2|} \rfloor = \tau + \lfloor \frac{\gamma' + \epsilon_1\epsilon_2}{n} \rfloor = \tau$$

hold.

Lemma 13 *(15) is $\tau+1$, if and only if we have $0 \le [\alpha m]_n < \alpha$.*

Proof. The equality $\lfloor \frac{\alpha m}{n} \rfloor - \lfloor \frac{\alpha(m-1)}{n} \rfloor = 1$ implies that there exists an integer n_0 satisfying $\frac{\alpha(m-1)}{n} < n_0 \le \frac{\alpha m}{n}$. Therefore equivalently $0 \le [\alpha m]_n < \alpha$ holds. $\square$

Since we have

$$\begin{aligned} \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} &= \frac{p + (\epsilon_1\epsilon_2 k + 1)n_1}{ak_1} \\ &= \frac{ak_1^2 + (\epsilon_1\epsilon_2 k_1 + 1)(nn_1 + \epsilon_2)}{ank_1} \\ &= \tau + \epsilon_1\epsilon_2 + \frac{\gamma' - \epsilon_1\epsilon_2 n'}{n} + \frac{n-n'}{nk_1}, \end{aligned} \tag{16}$$

$\frac{n-n'}{nk_1} < \frac{1}{k_1} \le \frac{1}{4n}$, and $\gamma' - \epsilon_1\epsilon_2 n' = (\gamma')^{-1} n' \ne 0 \ (n)$, we have

$$\lfloor \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} \rfloor = \tau + \epsilon_1\epsilon_2 + \lfloor \frac{\gamma' - \epsilon_1\epsilon_2 n'}{n} \rfloor \le \tau + 1.$$

$\square$

Lemma 14 *For any $0 < i < a$ we have*

$$\frac{[-\epsilon_2 n_i q_1]_p}{k_1} = i\tau + \frac{i\gamma' + \epsilon_1\epsilon_2 n_i'}{n} + \frac{n_i'}{nk_1}$$

and

$$\lfloor \frac{[-\epsilon_2 n_i q_1]_p}{k_1} \rfloor = i\tau + \epsilon_1\epsilon_2 + \lfloor \frac{i\gamma' - \epsilon_1\epsilon_2 [n'i]_n}{n} \rfloor.$$

Proof

$$\begin{aligned} \frac{[-\epsilon_2 n_i q_1]_p}{k_1} &= \frac{ip + (\epsilon_1\epsilon_2 k_1 + 1)n_i}{ak_1} \\ &= \frac{aik_1^2 + (\epsilon_1\epsilon_2 k_1 + 1)(nn_i + i\epsilon_2)}{ank_1} \\ &= i\tau + \frac{i\gamma' + \epsilon_1\epsilon_2 n_i'}{n} = i\tau + \frac{i\gamma' + \epsilon_1\epsilon_2 n_i'}{n} + \frac{n_i'}{nk_1} \end{aligned}$$

Here $\frac{n_i'}{nk_1} \le \frac{n}{nk_1} \le \frac{1}{2n}$ and $i\gamma' + \epsilon_1\epsilon_2 n_i' = ia^{-1}(\gamma')^{-1}(a(\gamma')^2 + \epsilon_1\gamma') = ia^{-1}(\gamma')^{-1}(-\epsilon_2) \ne 0(n)$ hold. Hence we have $\lfloor \frac{i\gamma' + \epsilon_1\epsilon_2 n_i'}{n} + \frac{n_i'}{nk_1} \rfloor = \lfloor \frac{i\gamma' + \epsilon_1\epsilon_2 n_i'}{n} \rfloor$. Since $n_i' = [n_1' i]_n = n - [n'i]_n$ holds, we get the formula as above. $\square$

Lemma 15 *For any $0 < i < a$ the integer l_i satisfying*

$$\lfloor \frac{pl_i}{|q_2|} \rfloor = [-\epsilon_2 n_i q_1]_p + \begin{cases} 0 & \epsilon_1\epsilon_2 = 1 \\ -1 & \epsilon_1\epsilon_2 = -1 \end{cases}$$

is $l_i = i(n\tau + \gamma' - \epsilon_1\epsilon_2) + \epsilon_1\epsilon_2 n'_i$.

5.4.1 The $\gamma' + \epsilon_1\epsilon_2(n - n') \geq n$ case.

Suppose that $\gamma' + \epsilon_1\epsilon_2(n-n') \geq n$ holds. Since $\gamma' \leq n-2$, we have $\epsilon_1\epsilon_2 = 1$. Therefore $\gamma' > n'$ and $\lfloor \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} \rfloor = \tau + 1$ hold.

If $\epsilon_1 = -1$, then we can find a forbidden subsequence as in the left of Figure 28, which this figure is a part of the global view of $A(i + jk_1)$. Hence none of data is realized by lens space surgery over S^3 or $\Sigma(2,3,5)$. Thus we assume that $\epsilon_1 = 1$. In this case we can get an admitted subsequence as the right in Figure 28.



Figure 28: The forbidden and admitted subsequences in the case of $\lfloor \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} \rfloor = \tau + 1$

Note that $A(x) = \tilde{a}_{\iota+\epsilon_1 x}$ for a constant ι independent of n_2. From Lemma 15 and the symmetry of Alexander polynomial we get Figure 29. In the case of $\epsilon_1\epsilon_2 = 1$, we have $l_1 = n(\tau + 1) - 1 + \gamma' - n'$. The block that $(i,j) = \partial^2_{lt}B$ satisfies $(i + jk_1) =$



Figure 29: The admitted subsequences $\epsilon_2 = 1$

$[-\epsilon_2 n_1 q_1]_p$ is $B_{\gamma'-n',0}$. Thus $\gamma' - n' = 2$ holds. From Lemma 10, we have

$$(n'+2)^2 - n'(n'+2) - n' = 0 \ (n),$$

thus we have $n' = n-4$. Regarding $a(n-n') = nn_1 + 1$ as mod$n$, we have $4a - 1 = 0 \ (n)$. As a result we can get

$$\begin{cases} a = \frac{un+1}{4} & n_1 = u \\ n' = n-4, & \gamma' = n-2 \end{cases}$$

for some integer u.

On the other hand we have

$$-(-\bar{g} + (a-1)k_1 + 1) = -\bar{g} + (a-1)k_1 - ([-\epsilon_2 n_1 q_1]_p + k_1)k_2 + 1.$$

Hence $n_1 = a-1$ and $n_{a-1} = 1$.

Thus $u = a-1$ and we have

$$u(n-4) = 3.$$

Solving this equation, we get $(a, n, \gamma') = (2, 7, 5), (4, 5, 3)$, that is,

$$(p, k_1) = (14\tau^2 + 21\tau + 8, 7\tau + 5), (20\tau^2 + 25\tau + 8, 5\tau + 3).$$

Therefore these families are realized by A_1 and A_2 of Table 2.

5.4.2 The $\gamma' + \epsilon_1\epsilon_2(n-n') < 0$ case.

Suppose that $\gamma' + \epsilon_1\epsilon_2(n-n') < 0$ holds. Since $\gamma' \leq n-2$, we have $\epsilon_1\epsilon_2 = -1$. Then

$$\lfloor \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} \rfloor = \tau - 1$$

holds.

In the case of $\epsilon_1 = 1$ we get the left of Figure 30 in which there is a forbidden subsequence as indicated. Then if we are to find lens space surgery on S^3 or $\Sigma(2,3,5)$, then ϵ_1 must be -1.

In the case of $\epsilon_1 = -1$ we get the right of Figure 30, so that we can find the admitted subsequence in the box indicated. From the symmetry of Alexander polynomial $B_{-1,0}$ and $B_{0,0}$ are symmetrical about $(\tau+1, -\frac{\tau+3}{2})$ as in Figure 31. In the other words, if one rotate the global view by 180° about $(\tau+1, -\frac{\tau+3}{2})$, then one get the same global view. From Lemma 15 the block $B_{0,\star}$ that the second boundary $(i,j) = \partial^2_{tl} B_{0,\star}$ satisfies $i + jk_1 = [-\epsilon_2 n_{a-1} q_1]_p$ is $B_{n-n'-\gamma',1}$.

Hence we have $n - n' - \gamma' = 2$. From Lemma 10, in the same way

$$\begin{cases} a = \frac{un+1}{4} & n_1 = u \\ n' = n-4 & \gamma' = 2. \end{cases}$$

On the other hand we have

$$-(-\bar{g} + (-a-1)k_1 + 1) = -\bar{g} + (-a-1)k_1 - ([-\epsilon_2 n_{a-1} q_1] + 3 - 2k_1)k_2 + 1.$$

Thus we have $n_{a-1} = 1$ and $a_1 = a-1$. Hence we have $(n-4)u = 3$. Then $(u, n) = (1, 7), (3, 5)$. In the former case

$$(p, k_1) = (14\tau^2 + 7\tau + 1, 7\tau + 2)$$

In the latter case

$$(p, k_1) = (20\tau^2 + 15\tau + 3, 5\tau + 2)$$

In this next section we treat the case of $0 \leq \gamma' + \epsilon_1\epsilon_2(n-n') < n$.


j
i
0
−1
0
τ
−2
−τ
0
0
0
j
i
0
−1
1
τ + 1
−2
−τ
0
0
0

Figure 30: The forbidden and admitted subsequence in the case of $\gamma' + \epsilon_1\epsilon_2(n - n') < 0$.


j
i
0
−1
1
τ + 1
−2
−τ
0
0
0
0
0
$B_{n-n'-\gamma',-1}$

Figure 31: The admitted subsequence in the case of $\epsilon_1 = -1$.

5.5 The $2 \leq \gamma' \leq n-2$ case (ii).

Suppose that $0 \leq \gamma' + \epsilon_1\epsilon_2(n-n') < n$ holds. We claim the following Lemmas.

Lemma 16 *For i with $1 \leq i \leq a-1$ we have*

$$\lfloor \frac{p-[-\epsilon_2 n_{a-i}q_1]_p}{k_1} \rfloor = i\tau + \lfloor \frac{\gamma' i - \epsilon_1\epsilon_2(n-n'_i)}{n} \rfloor = i\tau + \lfloor \frac{\gamma' i - \epsilon_1\epsilon_2[n'i]_n}{n} \rfloor.$$

Proof Calculate as follows:

$$\begin{aligned} \frac{p-[-\epsilon_2 n_{a-i}q_1]_p}{k_1} &= \frac{ik_1}{n} + \frac{-\epsilon_1\epsilon_2(n'_{a-i}-\epsilon_2)}{n} + \frac{-n'_{a-i}+\epsilon_2}{nk_1} \\ &= \frac{ik_1}{n} + \frac{-\epsilon_1\epsilon_2(n-n'_i)}{n} - \frac{n-n'_i}{nk_1} \\ &= i\tau + \frac{i\gamma' - \epsilon_1\epsilon_2(n-n'_i)}{n} - \frac{n-n'_i}{nk_1}. \end{aligned}$$

Here since we have

$$i\gamma' - \epsilon_1\epsilon_2(n-n'_i) = i\gamma' + \epsilon_1\epsilon_2 n'_i = a^{-1}i(a\gamma' + \epsilon_1) = a^{-1}i(-\epsilon_2(\gamma')^{-1}) \neq 0 \quad (n)$$

Thus since $\frac{i\gamma'-\epsilon_1\epsilon_2(n-n'_i)}{n} \notin \mathbb{Z}$ and $\frac{n-n'_i}{nk_1} < \frac{1}{k_1} < \frac{1}{2n}$, we have

$$\lfloor \frac{i\gamma' - \epsilon_1\epsilon_2(n-n'_i)}{n} - \frac{n-n'_i}{nk_1} \rfloor = \lfloor \frac{i\gamma' - \epsilon_1\epsilon_2(n-n'_i)}{n} \rfloor.$$

□

Here we put $n_0 = 0$.

Lemma 17 *Let l'_i be the integer satisfying*

$$\lfloor \frac{(|q_2| - l'_i)p}{|q_2|} \rfloor = [-\epsilon_2 n_{a-i} q_1]_p.$$

Then l'_i is

$$in\tau + i\gamma' - \epsilon_1\epsilon_2(n - n'_i + i) = ik_1 - \epsilon_1\epsilon_2(n - n'_i + i).$$

Proof. Since

$$\lfloor \frac{(|q_2| - l'_i)p}{|q_2|} \rfloor = p - \lfloor \frac{l'_i p}{|q_2|} \rfloor - 1,$$

by easy calculation we get

$$\begin{aligned} \frac{(ik_1 - \epsilon_1\epsilon_2(n-n'_i+i))p}{|q_2|} &= p - \frac{(a-i)p + (\epsilon_1\epsilon_2 k_1 + 1)n_{a-i}}{a} - \frac{\epsilon_1\epsilon_2 n_{a-i}}{|q_2|} \\ &= p - [-\epsilon_2 n_{a-i}q_1]_p - \frac{\epsilon_1\epsilon_2 n_{a-i}}{|q_2|}. \end{aligned}$$

Thus putting $l'_i = ik_1 - \epsilon_1\epsilon_2(n - n'_i + i)$, we get

$$\lfloor \frac{(|q_2| - l'_i)p}{|q_2|} \rfloor = [-\epsilon_2 n_{a-i}q_1]_p + \begin{cases} 0 & \epsilon_1\epsilon_2 = 1 \\ -1 & \epsilon_1\epsilon_2 = -1. \end{cases}$$

□

We put $n' = [-\epsilon_2 a^{-1}]_n$. Then we have

$$l'_i = i(n\tau + \gamma' - \epsilon_1\epsilon_2) - \epsilon_1\epsilon_2[n'i]_n,$$

for $n - n'_i = [-\epsilon_2 a^{-1} i]_n$ and $n' = [-\epsilon_2 a^{-1}]_n$.

5.5.1 The $a = 2$ case

Lemma 18 *Let (p, k_1) be a data realized by lens space surgery of S^3 or $\Sigma(2,3,5)$. In the $a = 2$ case we get IX, X or E_i.*

Here we will classify the case where $n_1 = n_{a-1} \Leftrightarrow a = 2$ holds. We may assume that $n' \geq \gamma' + 2$ holds.

Thus we have $n_1 = n_{a-1} = 1,\ \gamma' = \frac{\gamma - \epsilon_2}{4},\ n' = \frac{n-\epsilon_2}{2}$. Hence from $\gamma' \leq n' - 2$,

$$\gamma \leq 2n - \epsilon_2 - 8$$

holds.

Suppose that $\epsilon_2 = 1$. If $\gamma = 2n - 9$, then from the integrality of p, we get $(\gamma, n) = (13, 11)$. This case $(p, k) = (22\tau^2 + 13\tau + 2, 11\tau + 3)$ is realized by (X).

Suppose that $\epsilon_2 = -1$. If $\gamma = 2n - 7$, then the from integrality of p, we get $(\gamma, n) = (3, 5)$, hence $\gamma' = 1$. This family before we classified.

We may assume that $\gamma < 2n - \epsilon_2 - 12$ holds.

Now since $\lfloor \frac{n}{\alpha} \rfloor \geq n' - \gamma'$ holds, using $n' - \gamma' = \frac{2n-\gamma-\epsilon_2}{4}$, we have

$$\frac{4n}{\gamma - \epsilon_2 + 4} > \frac{2n - \gamma - 1}{4} \Leftrightarrow n < \frac{(\gamma+1)(\gamma - \epsilon_2 + 4)}{2(\gamma - \epsilon_2 - 4)}.$$

Hence

$$\frac{\gamma + 12 + \epsilon_2}{2} < \frac{(\gamma+1)(\gamma - \epsilon_2 + 4)}{2(\gamma - \epsilon_2 - 4)}$$

Thus we have

$$\gamma < \frac{53 + 15\epsilon_2}{3 + \epsilon_2}.$$

Hence if $\epsilon_2 = 1$, then $\gamma \leq 16$, and if $\epsilon_2 = -1$, then $\gamma \leq 18$.

If $\epsilon_2 = 1$, then by easy calculation, the possibility is $(\gamma, n) = (9, 11)$ only. Thus $(p, k_1) = (22\tau^2 + 9\tau + 1, 11\tau + 2)$, and each of this family is realized by (IX).

If $\epsilon_2 = -1$, then by easy calculation, the possibility is $(\gamma, n) = (15, 27)$ only. This case does not satisfy $\lfloor \frac{n}{\alpha} \rfloor \geq n' - \gamma'$.

Now suppose that $\lfloor \frac{n}{\alpha} \rfloor \leq n' - \gamma'$ holds. Thus $\epsilon_2 = -1$ holds and from the symmetry of Alexander polynomial

$$n' - \gamma' = 2\lfloor \frac{n}{\alpha} \rfloor \Leftrightarrow \frac{n' - \gamma'}{2} \leq \frac{n}{\gamma' + 1} < \frac{n' - \gamma'}{2} + 1.$$

If $\gamma > 3$, then we have

$$\gamma' + 2 + \frac{2}{\gamma' - 3} < n' \leq \gamma' + 4 + \frac{10}{\gamma' - 3},$$

in particular $4 \leq n' - \gamma'$.

We assume that $n' = \gamma' + 4$. Then from Lemma 10 we get $n|27$, namely $n = 3, 9, 27$. The possibilities of them are $n = 27, 9$. If $n = 27$, then $n' = 14,\ \gamma' = 10$, and $\gamma = 39$. Therefore the initial data is

$$(p, k_1) = (54\tau^2 + 39\tau + 7, 27\tau + 10),$$

and this family is E_2. If $n = 9$, then $n' = 5,\ \gamma' = 1$. This case is classified before.

We assume that $n' \geq \gamma' + 6$. In the inequality above $3 \leq \gamma' \leq 8$ holds. All the possibilities are $(\gamma', n, n') = (8, 27, 14), (7, 25, 13), (6, 23, 12), (6, 25, 13), (5, 21, 11), (5, 23, 12), (5, 25, 13), (5, 27, 14), (4, 21, 11), (4, 23, 12), (4, 25, 13), (4, 27, 14), (4, 29, 15), (4, 31, 16),$



j
0
τ
$(\lfloor \frac{n}{\alpha} \rfloor - 1)\tau$
$\lfloor \frac{n}{\alpha} \rfloor\tau$
i
0
$-\tau+2$
$-\tau+1$
$B_{0,-1}$
$B_{1,-1}$
$B_{\lfloor \frac{n}{\alpha} \rfloor-1,-1}$
$B_{\lfloor \frac{n}{\alpha} \rfloor,-1}$

Figure 32: An admitted subsequence in the case of $a = 2$

(4, 33, 17) or (4, 35, 18). The data which satisfies Lemma 10 is (4, 27, 14) only. This case

$$(p, k_1) = (54\tau^2 + 15\tau + 1, 27\tau + 4)$$

is class E_2.

If $\gamma' = 3$, then $9 - 3n' - n' = 9 - 2(n+1) = -2n + 7 = 7 = 0\ (n)$, then $n = 7$ holds. Thus $n' = 4, \gamma' = 2$ holds. This does not satisfy $an = nn_{a-1} + 1$. ☐

Now we may assume that $a \geq 3$.

5.5.2 The $\epsilon_1\epsilon_2 = 1$ case.

First we suppose that $n' - \gamma' = 1$ holds. From Lemma 10 $n | 2\gamma' + 1$ hence

$$[-\epsilon_2 n_1 q_1]_p = k_1(\tau + 1) + 1 - \tau + \frac{2\gamma' + 1}{n}. \tag{17}$$

Therefore $n | 2\gamma' + 1$ holds. Now since

$$2\gamma' + 1 \leq 2(n - 2) + 1 = 2n - 3,$$

$2\gamma' + 1 = n$ must be true as $\mathbb{Z}$. Thus we have

$$\gamma' = \frac{n-1}{2},\ n' = \frac{n+1}{2}.$$

Taking $(n+1)a = (nn_{a-1} - \epsilon_2)2$ as $\bmod n$, we get $n | a + 2\epsilon_2$. Let u be the integer with $a = un - 2\epsilon_2$. Hence

$$n_{a-1} = \frac{a+u}{2},\ u_1 = \frac{a-u}{2}.$$

Therefore

$$(p, k_1) = (uk_1^2 - \epsilon_2\tau'(k_1 - 1), \frac{n\tau' - 1}{2}),$$

where $\tau' = 2\tau + 1$. These families are realized by (IV_-). (Put $-2\epsilon_2\tau' = J$.)

Hence we may assume that $n' \geq \gamma' + 2$.

Lemma 19 *Let (p, k_1) be a data with $\epsilon_1\epsilon_2 = 1$. Suppose that $0 < n' - \gamma' < n$ and $n' \geq \gamma' + 2$.*

1. *If $\epsilon_1 = 1$, then $\lfloor \frac{n}{\alpha} \rfloor = n' - \gamma'$ holds.*

2. *If $\epsilon_1 = -1$, then $\lfloor \frac{n}{\alpha} \rfloor = n' - \gamma'$ or $\lfloor \frac{2n}{\alpha} \rfloor = n' - \gamma'$ holds.*

Proof Suppose that $(\epsilon_1, \epsilon_2) = (1, 1)$. If $\lfloor \frac{n}{\alpha} \rfloor > n' - \gamma'$, then from the global view Figure 33, the indicated subsequence is the forbidden subsequence. Then we deduce that $\lfloor \frac{n}{\alpha} \rfloor \leq n' - \gamma'$, namely $\lfloor \frac{n}{\alpha} \rfloor = n' - \gamma'$ since $B(\lfloor \frac{n}{\alpha} \rfloor, -1)$ is the leftmost block with the top width $\tau + 1$ in $B(\star, -1)$ satisfying $\star \geq 0$.


j
0
τ
$\lfloor \frac{n}{\alpha} \rfloor \tau$
$(\lfloor \frac{n}{\alpha} \rfloor + 1)\tau$
i
0
$-\tau + 2$
$-\tau + 1$
$B_{0,-1}$
$B_{\lfloor \frac{n}{\alpha} \rfloor - 1, -1}$
$B_{\lfloor \frac{n}{\alpha} \rfloor, -1}$

Figure 33: A forbidden or admitted subsequence in the case of $\tau = 5$

Suppose that $(\epsilon_1, \epsilon_2) = (-1, -1)$. If $\lfloor \frac{n}{\alpha} \rfloor > n' - \gamma'$, in the same way we can find an admitted subsequence as in Figure 33. If $\lfloor \frac{2n}{\alpha} \rfloor > n' - \gamma'$, then we can find two admitted subsequences. Hence we have $\lfloor \frac{2n}{\alpha} \rfloor \leq n' - \gamma'$. Namely we have $\lfloor \frac{n}{\alpha} \rfloor = n' - \gamma'$ or $\lfloor \frac{2n}{\alpha} \rfloor = n' - \gamma'$ $\square$

Lemma 20 *Let (p, k_1) be a data realized by S^3 or $\Sigma(2, 3, 5)$. Suppose that $\epsilon_1 = -1$ and $\lfloor \frac{2n}{\alpha} \rfloor = n' - \gamma'$, and $a \geq 3$ holds. Then $2\lfloor \frac{n}{\alpha} \rfloor = n' - \gamma'$ holds and (p, k_1) is realized by type C_i or D_i for $i = 1, 2$.*

Proof. From Section 5.5, $0 < n' - \gamma' \leq n$ and Lemma 16 we have $\lfloor \frac{p - [-\epsilon_2 n_{a-1} q_1]_p}{k_1} \rfloor = \tau + \lfloor \frac{\gamma' - n'}{n} \rfloor = \tau - 1$. Thus the height of $B_{0,-1}, \cdots, B_{n'-\gamma'-1,-1}$ are all τ. From


j
τ
$i\tau$
$(i+1)\tau$
i
O
$-\tau + 2$
$-\tau + 1$
0
$B_{0,-1}$
$B_{i,-1}$
$B_{n'-\gamma',-1}$
symmetry

Figure 34: A forbidden or admitted subsequence for Lemma 20 and the symmetry.

Lemma 13 and the assumption, $B_{\lfloor \frac{n}{\alpha} \rfloor, -1}$ is top width $\tau + 1$. Then $n' - \gamma' \geq 2$ holds.

We define a positive integer i to be $n' - \gamma' = 2i$. From the top width of $B_{j,-1}$ $(0 \leq j < i)$ is τ and the top width of $B_{i,-1}$ is $\tau + 1$, we have

$$\alpha \leq [\alpha j]_n \ (0 \leq j < i), \ \ 0 \leq [\alpha i]_n < \alpha$$

This condition is equivalently

$$\gamma' + 1 \leq [(\gamma' + 1)j]_n \ (0 \leq j < i), \ \ 0 \leq [(\gamma' + 1)i]_n < \gamma' + 1, \tag{18}$$

for $0 < \frac{aj}{|q_2|} < 1$ holds. The condition (18) implies

$$n < i(\gamma' + 1) < n + \gamma' + 1 < (i + 1)(\gamma' + 1).$$

Thus the inequality

$$\frac{n - i}{i} < \gamma' < \frac{n + 1 - i}{i - 1} \tag{19}$$

From the integrality condition and $\gamma' + 2i = n'$, we have $(2i + 1)\gamma' = -2i$. Thus for some positive integer u' we have $\gamma' = \frac{u'n - 2i}{2i + 1}$. The integer u' satisfying (19) is 2 only. Thus we have

$$\begin{cases} a = \frac{un + 2i + 1}{4i^2} & n_1 = \frac{(2i-1)a - u}{2i+1} \\ n' = \frac{2n + 4i^2}{2i+1} & \gamma' = \frac{2n - 2i}{2i+1}. \end{cases}$$

The symmetry of Alexander polynomial induces symmetry of the global view as indicated in Figure 34. From the symmetry of the global view we have

$$-(-\bar{g} + (a - 1)k_1 + 1) = -\bar{g} + (a - 1)k_1 - (-\epsilon_2 n_{a-1} q_1 + 2 + k_1)k_2 + 1,$$

thus we have $n_1 = 1$. Thus we have

$$((2i - 1)n - 4i^2)u = 8i^3 + 1. \tag{20}$$

In the case of $i = 1$, (20) is $u(n - 4) = 9$, thus $(u, n) = (1, 13), (9, 5), (3, 7)$. The data satisfying integral condition of n' are

$$(p, k_1) = (52\tau^2 + 63\tau + 19, 13\tau + 8), (42\tau^2 + 47\tau + 13, 7\tau + 4).$$

In the case of $i = 2$, (20) is $u(3n - 16) = 65$, thus $(u, n) = (1, 27), (13, 7)$. Thus we have

$$(p, k_1) = (54\tau^2 + 39\tau + 7, 27\tau + 10), (42\tau^2 + 23\tau + 3, 7\tau + 2).$$

The former case is inconsistent with the assumption $a \geq 3$.

In the case of $i = 3$ the equation (20) is $u(5n - 36) = 217$, thus $(u, n) = (1, 217), (217, 1), (7, 31), (31, 7)$. The data satisfying integral condition of n' is $(u, n) = (7, 31)$. The case $(7, 31)$ fails to integrality of a.

In the case of $i = 4$ the equation (20) is $u(7n - 64) = 513$, thus $(u, n) = (19, 13)$. Thus we have

$$(p, k_1) = (52\tau^2 + 15\tau + 1, 13\tau + 2).$$

In the case of $i = 5$ the equation (20) is $u(9n - 100) = 1001$, thus $(u, n) = (19, 27)$. This is inconsistent with the assumption $a \geq 3$.

Suppose that $i \geq 6$. From

$$n' - \gamma' = 2\lfloor \frac{n}{\alpha} \rfloor < 2\frac{n}{\gamma' + 1}$$

we can get

$$n' < \frac{n_{a-1}\gamma'(\gamma'+1)-2}{n_{a-1}(\gamma'+1)-2a}.$$

Hence we have

$$\gamma'+12 < \frac{n_{a-1}\gamma'(\gamma'+1)-2}{n_{a-1}(\gamma'+1)-2a} \Rightarrow \quad \gamma' < \frac{12a-1}{5n_{a-1}-n_1}-1 = \frac{7a+5}{5a-6} \leq \frac{26}{9}.$$

Hence $\gamma'=2$ holds.

In the case of $\gamma'=2$, $n=3i+1$ and

$$u((2i-1)(3i+1)-4i^2)=(2i+1)(4i^2-2i+1) \Leftrightarrow u(2i+1)(i-1)=(i-1)(4i+2)+3.$$

Then $i-1|3$ Hence $i=2,4$. This case is classified before. Therefore the appearing cases are all realized by type C_i, D_i. $\square$

Suppose that $n'-\gamma'=\lfloor\frac{n}{\alpha}\rfloor$ holds.

Lemma 21 *Let (p,k_1) be a data with $\epsilon_1\epsilon_2=1$ and $\lfloor\frac{n}{\alpha}\rfloor = n'-\gamma' \geq 2$. Then we have $0<n'<\frac{n}{2}$.*

Proof Suppose that $\frac{n}{2}<n'<n$ holds. In the case of $n'=\gamma'+i \quad (i\geq 2)$ we have $(n'-i)^2-n'(n'-i)-n'=0 \ (n)$, $(i+1)n'=i^2 \ (n)$. From $\alpha \leq [\alpha(n(-\tau+1)+j+1)]_n < n \quad (j=1,2,\cdots,i-1)$ and $\alpha\leq[\alpha(n(-\tau+1)+i+1)]_n<n$, we have

$$\frac{n+i(i+1)}{i+1}-1 \leq n' < \frac{n+i^2}{i}-1.$$

Thus we have $n'=\frac{n+i^2}{i+1}$. Then we have

$$\begin{cases} a=\frac{un-\epsilon_2(i+1)}{i^2} & n_1=\frac{ia-u}{i+1} \\ n'=\frac{n+i^2}{i+1} & \gamma'=\frac{n-i}{i+1}. \end{cases}$$

From $\frac{n}{2}<\frac{n+i^2}{i+1}$ we have $n<2(i+1)+\frac{2}{i-1}$. If $i\geq 3$, we have $n<2i+3$. Since $i=\lfloor\frac{n}{\alpha}\rfloor<\frac{n}{3}<\frac{2i+3}{3}$, we have $i<3$. This is a contradiction. If $i=2$, then we have $\gamma'=\frac{n-2}{3}$ and $n<8$. The possibility is $n=5$ only since the integrality of γ'. In this case $a=\frac{5u-3\epsilon_2}{4}$, $n_1=\frac{2a-u}{3}$, $n'=3$ and $\gamma'=1$. This case is already classified before. Therefore we have $0<n'<\frac{n}{2}$. $\square$

Lemma 21 holds even in the case of $\epsilon_1\epsilon_2=-1$ replacing γ' and α with $n-\gamma'$ and $n-\alpha$. The proof is skipped because the process is similar to Lemma 21.

Lemma 22 *Let (p,k_1) be a stable data. If $a\geq 3$, $0<n'-\gamma'<n$ $\epsilon_1=1$ and $0<n'<\frac{n}{2}$, then none of data satisfying this condition is realized by S^3 or $\Sigma(2,3,5)$.*

Proof From Lemma 16, we get

$$\lfloor\frac{p-[-\epsilon_2 n_{a-2}n_1q_1]_p}{k_1}\rfloor = 2\tau+\lfloor\frac{2\gamma'-2n'}{n}\rfloor = 2\tau-\lfloor\frac{2n'-2\gamma'}{n}\rfloor-1=2\tau-1.$$

Thus we can find a forbidden subsequence from the Figure 35. $\square$

Then Lemma 16 $\lfloor\frac{p-[-\epsilon_2 n_{a-2}q_1]_p}{k_1}\rfloor = 2\tau+\lfloor\frac{2\gamma'-2n'}{n}\rfloor = 2\tau-1$ holds, because $n'-\gamma'<\frac{n}{2}$.

Lemma 23 *Let (p,k_1) be a data. Suppose that $\epsilon_1=-1$. If $n'-\gamma'=\lfloor\frac{n}{\alpha}\rfloor$, then the data (p,k_1) is either of $F_1, F_2, G_1, G_2, H_1,$ or H_2.*



Figure 35: The place in which the heights of blocks change.($0 < n' < \frac{n}{2}$)

Proof. The global view is as in Figure 35 since $0 < n' < \frac{n}{2}$ and Lemma 16. From this view we have

$$-(-\bar{g}+(a-1)k_1+1) = -\bar{g}+(a-1)k_1-(-\epsilon_2 n_{a-2}q_1+2+k_1)k_2+1,$$

$n_2 = 1$, hence $n_1 = \frac{a+1}{2}$.

Putting $n'-\gamma' = i$, we have

$$\begin{cases} a = \frac{un+i+1}{i^2} & n_1 = \frac{ia-u}{i+1} \\ n' = \frac{n+i^2}{i+1} & \gamma' = \frac{n-i}{i+1}. \end{cases}$$

Since $a = \frac{2u+i+1}{i-1}$,

$$((i-1)n-2i^2)u = (i+1)(i^2-i+1)$$

If $i \leq 5$ holds, the possibility is $(i,u,n,a,n',\gamma') = (2,1,17,5,7,5), (2,3,11,9,5,3), (3,7,11,9,5,2),$ $(3,1,23,3,8,5)$, $(5,7,17,5,7,2), (5,3,23,3,8,3)$

$$(p,k_1) = (85\tau^2+49\tau+7, 17\tau+5), (99\tau^2+53\tau+7, 11\tau+3)$$

$$(p,k_1) = (99\tau^2+35\tau+3, 11\tau+2), (69\tau^2+29\tau+3, 23\tau+5)$$

$$(p,k_1) = (85\tau^2+19\tau+1, 17\tau+2), (69\tau^2+17\tau+1, 23\tau+3).$$

If $i \geq 6$ holds, $\gamma' < \frac{3a-2}{2a+3} < \frac{3}{2}$ holds. Since the assumption is $\gamma' \geq 2$, we can get to the assertion. $\square$

5.5.3 The $\epsilon_1\epsilon_2 = -1$ case.

We assume that $0 < \gamma'-n+n' < n$ and $\epsilon_1\epsilon_2 = -1$.

We consider $\gamma'-n+n' = 1$. From integral condition $2n'-1 = 0$ (n) holds. Then we have $n' = \frac{n+1}{2}$. Thus we have $\gamma' = \frac{n+1}{2} = n'$. We have $k_1 = n\tau + \frac{n+1}{2}$. Taking $2a(n-n') = 2nn_1 + 2\epsilon_2$ as $\mathbb{Z}/p\mathbb{Z}$, we have $a = -2\epsilon_2$ (n). For some integer u we have $a = un - 2\epsilon_2$. Then we have $(p,k_1) = (uk_1^2 - \epsilon_2(2\tau+1)(k_1+1), n\tau + \frac{n+1}{2})$. This case is realized by type (IV$_+$).



Figure 36:

We may assume that $\gamma'-n+n'\geq 2$ holds. Hence $\lfloor\frac{[-\epsilon_2 n_1 q_1]_p}{k_1}\rfloor=\tau-1+\lfloor\frac{n'+\gamma'}{n}\rfloor=\tau$ holds.

Now we assume that $\lfloor\frac{(n\tau+2)p}{|q_2|}\rfloor-\lfloor\frac{(n\tau+1)p}{|q_2|}\rfloor=\tau$. Equivalently $\alpha\leq[\alpha(n\tau+2)]_n<n$ holds. Here we have $[\alpha(n\tau+2)]_n=[2(\gamma'-1)+\frac{an\tau+2a}{|q_2|}]_n$ and

$$\frac{an\tau+2a}{|q_2|}=1-\frac{a\gamma'-a+\epsilon_1}{|q_2|}<1.$$

Thus $\alpha<[2\gamma'-2+\frac{an\tau+2a}{|q_2|}]_n$ holds and $2\gamma'-2<n$ holds. As a result $\gamma'<\frac{n+2}{2}$ holds. We put $\gamma''=n-\gamma'$ and $\alpha'=\gamma''+1-\frac{a}{|q_2|}=n-\alpha$.

Lemma 24 *Let (p,k_1) be a data with $\epsilon_1\epsilon_2=-1$, $0<n'-\gamma''<n$, and $n'\geq\gamma''+2$. Then the possibilities are either of the following.*

1. *If $\epsilon_1=-1$, then $\lfloor\frac{n}{\alpha'}\rfloor=n'-\gamma''$.*
2. *If $\epsilon_1=1$, then $\lfloor\frac{n}{\alpha'}\rfloor=n'-\gamma''$ or $\lfloor\frac{2n}{\alpha'}\rfloor=n'-\gamma''$.*

Proof. Suppose that $(\epsilon_1,\epsilon_2)=(-1,1)$. If $\lfloor\frac{n}{\alpha'}\rfloor<n'-\gamma''$, the global view of $A(i+jk_1)$ is as Figure 37. Here we must note that $B_{\lfloor\frac{n}{\alpha'}\rfloor-1,0}$ is the leftmost block with the top width τ in the blocks of $B_{*,0}$ $(*\geq 0)$. Therefore we can find a forbidden subsequence as indicated by the box. Thus we have $\lfloor\frac{n}{\alpha'}\rfloor\geq n'-\gamma''$. Since the block $B_{\lfloor\frac{n}{\alpha'}\rfloor,0}$ is the leftmost block the top width τ we have $\lfloor\frac{n}{\alpha'}\rfloor=n'-\gamma''$.

Suppose that $(\epsilon_1,\epsilon_2)=(1,-1)$. If $\lfloor\frac{2n}{\alpha'}\rfloor<n'-\gamma''$, the global view of $A(i+jk_1)$ is as Figure 38. In this time there are two admitted subsequences in this view. This is contradiction. Thus $\lfloor\frac{2n}{\alpha'}\rfloor\geq n'-\gamma''$ holds. Since the block $B_{n'-\gamma'',0}$ has the bottom width τ, $n'-\gamma''=\lfloor\frac{n}{\alpha'}\rfloor$ or $\lfloor\frac{2n}{\alpha'}\rfloor$. $\Box$

We first classify the case of $\lfloor\frac{2n}{\alpha'}\rfloor=n'-\gamma''$.

Lemma 25 *Suppose that $\epsilon_1=1$, $\lfloor\frac{2n}{\alpha'}\rfloor=n'-\gamma''$ and $a\geq 3$. Then the data is realized by one of C_i and D_i $(i=1,2)$.*


j
$B_{0,0}$
$B_{\lfloor \frac{n}{\alpha'} \rfloor,0}$
$\cdots$
$\cdots$
i

Figure 37: A global view of $(\epsilon_1, \epsilon_2) = (-1, 1)$


j
$B_{0,0}$
$B_{\lfloor \frac{n}{\alpha'} \rfloor,0}$
$B_{\lfloor \frac{2n}{\alpha'} \rfloor,0}$
$\cdots$
$\cdots$
$\cdots$
i

Figure 38: A global view of $(\epsilon_1, \epsilon_2) = (1, -1)$


j
τ
$[-\epsilon_2 n_1 q_1]_n$
0
$\cdots$
$\cdots$
i
-1
0
τ
$B_{i,0}$
$B_{\gamma'+n'-n,0}$

Figure 39: A global view of Lemma 25

Proof. From the assumption the global view of (p, k_1) is as in Figure 39. First for a positive integer j the equation $\frac{a(n\tau+1+j)}{|q_2|} = 1 - \frac{a\gamma'-aj+1}{|q_2|}$ holds. Furthermore we have

$$0 < \frac{a\gamma' - aj + 1}{|q_2|} < \frac{a(\gamma'-1)+1}{2an + a\gamma' - a + 1} = 1 - \frac{2an}{2an + a\gamma' - a + 1} < 1.$$

As a result we have $0 < \frac{a(n\tau+1+j)}{|q_2|} < 1$.

If

$$\lfloor \frac{(n\tau+1+j)p}{|q_2|} \rfloor - \lfloor \frac{(n\tau+j)p}{|q_2|} \rfloor = \tau + 1$$

holds, then from Lemma 13 $0 \leq [\alpha(n\tau+j+1)]_n = [(j+1)(\gamma'-1) + \frac{a(n\tau+1+j)}{|q_2|}]_n < \alpha$ holds. This condition is equivalent to the following:

$$\begin{aligned} &\Leftrightarrow \quad 0 \leq \left[(j+1)(\gamma'-1) + \frac{a(n\tau+i)}{|q_2|}\right]_n < \gamma' - 1 \\ &\Leftrightarrow \quad 0 \leq [(j+1)(\gamma'-1)]_n < \gamma' - 1. \end{aligned}$$

Now suppose that the global view is as Figure 39. Then we have $0 \leq [(j+1)(\gamma'-1)]_n < \gamma'-1$ for some integer with $0 < j < i$ and we have $\gamma'-1 \leq [(i+1)(\gamma'-1)]_n < n$. Therefore we have

$$(i-1)n \leq i(\gamma'-1) < (i-1)n + \gamma' - 1 < (i+1)(\gamma'-1) < in. \tag{21}$$

Hence

$$\frac{(i-1)n+i}{i} \leq \gamma' < \frac{in+i+1}{i+1}$$

holds.

In the Figure 39 we can find an admitted subsequence as indicated by a box in the figure. Furthermore from Lemma 15 the block $B_{\star,0}$ that the point $(i,j) = \partial^2_{lt} B_{\star,0}$ satisfies $i + jk_1 = [-\epsilon_2 n_1 q_1]_p$ and is the closest to the origin is $B_{\gamma'+n'-n,0}$. Now from the symmetry of Alexander polynomial we put $\gamma' + n' - n = 2i$. Hence (21) implies

$$\frac{n+2i^2+i-1}{i+1} < n' \leq \frac{n+2i^2-i}{i}.$$

From the integral condition $(2i+1)n' = 4i^2$ (n) and $4i^2 a = 2i+1$ (n) hold. Therefore $n' = \frac{2n+4i^2}{2i+1}$ and $a = \frac{un+2i+1}{4i^2}$ hold. Hence

$$\begin{cases} a = \frac{un+2i+1}{4i^2} & n_1 = \frac{(2i-1)a-u}{2i+1} \\ n' = \frac{2n+4i^2}{2i+1} & \gamma' = \frac{(2i-1)n+2i}{2i+1}. \end{cases} \tag{22}$$

Furthermore from the symmetry of Alexander polynomial

$$-(-\bar{g} + (-a-1)k_1 + 1) = -\bar{g} + (-a-1)k_1 + (-\epsilon_2 n_1 q_1 + 1 - 2k_1)k_2 + 1 \ (p).$$

Hence we have $n_1 = 1$. Using (22) and $n_1 = 1$

$$((2i-1)n - 4i^2)u = 8i^3 + 1.$$

In the case of $i = 1$ we get

$$(p, k_1) = (52\tau^2 + 41\tau + 8, 13\tau + 5), (42\tau^2 + 37\tau + 8, 7\tau + 3).$$

In the case of $i = 2$ we get

$$(p, k_1) = (54\tau^2 + 69\tau + 22, 27\tau + 17), (42\tau^2 + 61\tau + 22, 7\tau + 5).$$

The former case is inconsistent with $a \geq 3$. In the case of $i = 3$ there does not exist any suitable pair (u, n). In the case of $i = 4$ we get

$$(p, k_1) = (52\tau^2 + 89\tau + 38, 13\tau + 11).$$

In the case of $i = 5$ we get

$$(p, k_1) = (54\tau^2 + 93\tau + 40, 27\tau + 23).$$

This is inconsistent with $a \geq 3$.

If $i \geq 6$, then $\gamma' - n + n' = 2i \geq 12$ holds. From the form of the blocks, $\lfloor \frac{n}{n-\alpha} \rfloor = \frac{\gamma' - n + n'}{2}$ holds.

Thus if $\gamma'' \neq 2$ or $(\gamma'', a) \neq (3, 3)$ we have

$$n' < \frac{n_{a-1}(\gamma'')^2 - 2}{n_{a-1}\gamma'' - 2a},$$

where $\gamma'' = n - \gamma'$.

Hence we have

$$\gamma'' + 12 < \frac{n_{a-1}(\gamma'')^2 - 2}{n_{a-1}\gamma'' - 2a} \Leftrightarrow \gamma'' < \frac{12a - 1}{5n_{a-1} - n_1} = \frac{12a - 1}{5a - 6} \leq \frac{35}{9}.$$

Thus we have $\gamma'' = 3$ and $a \neq 3$.

If $\gamma'' = 3$, then from the integral condition we have $9 - 4n' = 0(n)$. Then we have $n' = \frac{n+9}{4}$ since $\frac{n'-3}{2} = \lfloor \frac{n}{4 - \frac{a}{|q_2|}} \rfloor < \frac{2n}{7}$. Thus we have $n = 8i + 3$ and $n' = 2i + 3$. Here since

$$i = \lfloor \frac{8i + 3}{4 - \frac{a}{|q_2|}} \rfloor \geq \lfloor \frac{8i + 3}{4} \rfloor = 2i$$

this is a contradiction.

If $\gamma'' = 2$, then from the integral condition we have $n' = \frac{2n+4}{3}$. Then we have $n = 3i + 1$, $(i - 1)u = 4i^2 - 2i + 1$ and $u = \frac{4i^2 - 2i + 1}{i-1} = 4i + 2 + \frac{3}{i-1}$ hold. Thus $i - 1 = 1, 3$ holds these are inconsistent with $i \leq 6$. □

Lemma 26 *If $0 < n' < \frac{n}{2}$ and $\lfloor \frac{n}{\alpha'} \rfloor = n' - \gamma''$ hold, then $\epsilon_1 = 1$ and the global view of a data (p, k_1) is Figure 40, the data is realized by one of type F_i, G_i and H_i for $i = 1, 2$.*

Proof From the assumption the global view is as Figure 40. For since from Lemma 14 $\lfloor \frac{[-\epsilon_2 n_1 q_1]_p}{k_1} \rfloor = \tau + \lfloor \frac{n' - \gamma''}{n} \rfloor = \tau$ and $\lfloor \frac{[-\epsilon_2 n_2 q_1]_p}{k_1} \rfloor = 2\tau - 1 + \lfloor \frac{2\gamma' + [2n']_n}{n} \rfloor = 2\tau + 1 + \lfloor \frac{2n' - 2\gamma''}{n} \rfloor = 2\tau + 1$, we have $H(B_{0,0}) = \tau + 3$ and $H(B_{0,1}) = \tau + 2$. The leftmost block $B_{*,0}$ in which $(i, j) = \partial^2_{lt} B_{*,0}$ satisfies $i + jk_1 = [-\epsilon_2 n_1 q_1]_p$ for positive integer $*$ is $B_{n' - \gamma'', 0}$. Namely $H(B_{j,0}) = \tau + 3$ $(0 \leq j < n' - \gamma'')$ and $H(B_{n' - \gamma'', 0}) = \tau + 2$. This is due to Lemma 15.

From the global view we can find a subsequence as indicated by the box in Figure 40. This subsequence is the forbidden subsequence when $\epsilon_1 = -1$, and the admitted subsequence when $\epsilon_1 = 1$. To realize lens space surgery over S^3 or $\Sigma(2, 3, 5)$ we must take $\epsilon_1 = 1$. Therefore 1. part in Lemma 24 does not occur as long as lens space surgery over S^3 or $\Sigma(2, 3, 5)$.



Figure 40: A forbidden or admitted subsequences with $\tau = 4$

From Lemma 25

$$\frac{n-i-1}{i+1} < \gamma'' \leq \frac{n-i}{i}$$

holds. We denote $n' - \gamma''$ by i.

From the integrality condition we have for some integer u, we get

$$\begin{cases} a = \frac{un+i+1}{i^2} & n_1 = \frac{ia-u}{i+1} \\ n' = \frac{n+i^2}{i+1} & \gamma'' = \frac{n-i}{i+1}. \end{cases}$$

Furthermore from the symmetry of Alexander polynomial we have

$$-(\bar{g} + (-a-1)k_1 + 1) = -\bar{g} + (-a-1)k_1 + (-\epsilon_2 n_2 q_1 + 1 - 2k_1)k_2 + 1$$

and $n_2 = 1$. Figure 41 illustrates the symmetry.

Thus we have $n_1 = \frac{a+1}{2}$. Hence

$$((i-1)n - 2i^2)u = i^3 + 1. \tag{23}$$

In the case of $i = 2$, the possibilities are $(n, u) = (9, 9), (11, 3), (17, 1)$. If $(n, u) = (9, 9)$, then $n' = \frac{13}{3} \notin \mathbb{Z}$, hence this is inconsistent. If $(n, u) = (11, 3), (17, 1)$, then

$$(p, k_1) = (99\tau^2 + 97\tau + 53, 11\tau + 8), (85\tau^2 + 121\tau + 43, 17\tau + 12).$$

In the case of $i = 3$, the possibilities are $(n, u) = (11, 7), (23, 1)$. Hence we have

$$(p, k_1) = (99\tau^2 + 163\tau + 67, 11\tau + 9), (69\tau^2 + 109\tau + 43, 23\tau + 18).$$

In the case of $i = 4$ the possibilities are $(n, u) = (11, 65), (15, 5)$. If $(n, u) = (11, 65)$, then $n' = \frac{27}{5} \notin \mathbb{Z}$, hence this is inconsistent. If $(n, u) = (15, 5)$, then $n' = \frac{31}{5} \notin \mathbb{Z}$, hence this is inconsistent as well.

In the case of $i = 5$, the possibilities are $(n, u) = (13, 63), (17, 7), (23, 3)$. If $(n, u) = (13, 63)$, then $n' = \frac{13}{2} \notin \mathbb{Z}$, hence this is inconsistent. If $(n, u) = (17, 7), (23, 3)$, then

$$(p, k_1) = (85\tau^2 + 151\tau + 67, 17\tau + 15), (69\tau^2 + 121\tau + 53, 23\tau + 20)$$



Figure 41: A forbidden or admitted subsequences with $\tau = 4$

If $i \geq 6$, then using $\lfloor \frac{n}{n-\alpha} \rfloor = n' - \gamma''$ we get

$$n' - \gamma'' < \frac{n}{\gamma'' + \frac{1}{2}},$$

so that we get

$$n' < \frac{n_{a-1}\gamma''(\gamma'' + \frac{1}{2}) - 1}{n_{a-1}(\gamma'' + \frac{1}{2}) - a}.$$

Hence we have $\gamma'' < \frac{9a+1}{2(2a-3)} = \frac{9}{4} + \frac{29}{4(2a-3)} \leq \frac{14}{3}$. Thus $\gamma'' \leq 4$ holds.

If $\gamma'' = 4$, then from the integrality condition $5n' = 16(n)$, then from $n' = \frac{n+16}{5} = \frac{n+i^2}{i+1}$ we have $n = 5i+4$. Using (23), we have $u = \frac{i^2-i+1}{3i-4}$. If $i = 3i_1$ for some positive integer i_1, then $u = \frac{9i_1^2-3i_1+1}{9i_1-4} = i_1 + \frac{i_1+1}{9i_1-4} \notin \mathbb{Z}$. If $i = 3i_1+1$ for some positive integer i_1, then $u = \frac{9i_1^2+3i_1+1}{9i_1-1} = i_1 + \frac{4i_1+1}{9i_1-1} \notin \mathbb{Z}$. Thus we cannot find any integer solution in (23), this implies that there is no lens space surgery on S^3 or $\Sigma(2,3,5)$ in this case.

If $\gamma'' = 3$, then from the integrality condition $4n' = 9(n)$, then from $n' = \frac{n+9}{4} = \frac{n+i^2}{i+1}$ we have $n = 4i+3$. Using (23), we have $u = \frac{i^2-i+1}{2i-3}$. If $i = 2i_1$ for some positive integer i_1, then $u = \frac{4i_1^2-2i_1+1}{2i_1-3} = i_1 + \frac{i_1+1}{4i_1-3} \notin \mathbb{Z}$. If $i = 2i_1+1$ for some positive integer i_1, then $u = \frac{4i_1^2+2i_1+1}{4i_1-1} = i_1 + \frac{3i_1+1}{4i_1-1}$. Thus we have $i_1 = 2$, $u = 3$, and $i = 5$ and this is inconsistent with $i \geq 6$. There is no lens space surgery on S^3 or $\Sigma(2,3,5)$ in this case as well.

If $\gamma'' = 2$ holds, then from the integrality condition $3n' = 4(n)$ and $n' = \frac{n+4}{3} = \frac{n+i^2}{i+1}$ we have $n = 3i+2$. Using (23), we have $u = \frac{i^2-i+1}{i-2} = i+1+\frac{3}{i-2}$. Thus we have $i-2 = 1,3$ these cases are inconsistent with $i \geq 6$. Therefore in the case of $i \geq 6$ we cannot find any lens space surgery over S^3 or $\Sigma(2,3,5)$. $\square$

In this point we can classify all stable data (p, k_1) with lens space surgery S^3 or $\Sigma(2,3,5)$. To prove that Berge's list is fully complete, we must argue unstable data: namely $\tau = 1$ or 0. This argument will be done in sequent papers.

Acknowledgments

This research is partially supported by the Grant-in-Aid for JSPS Fellow for Young Scientist (21-1458). The author is grateful for Professor M.Ue and T.Ohtsuki affiliated in Mathematics department and RIMS, Kyoto University respectively, who give me a lot of useful comments in a series of arguments. Furthermore a part of the main theorem was proved while my visit of Michigan State University consequently the author thanks for the hospitality of the institute and the acceptance of my visit by Professor Akbulut. Finally, the author deeply appreciates Professors T.Kadokami and Y.Yamada giving me the motivation for this research.

References

[1] Berge, *Some knots with surgeries yielding lens spaces* (unpublished manuscript, 1990)

[2] Fintushel and Stern, *Constructing lens spaces by surgery on knots* Math. Z. 175 (1980), no. 1, 33–51.

[3] Greene, *Donaldson's Theorem, Heegaard Floer Homology, and Results on Knots* PhD. thesis, Princeton Univ., (2009)

[4] T. Kadokami, and Y. Yamada, *A deformation of the Alexander polynomials of knots yielding lens spaces* Bull. Austral. Math. Soc. 75 (2007), no. 1, 75–89.

[5] P. Ozsváth and Z. Szabó, *On knot Floer homology and lens space surgery*, Topology, 44 (2005), no.6, 1281-1300.

[6] P. Ozsváth and Z. Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. 173 (2003), no. 2, 179–261.

[7] M. Tange, *Ozsváth-Szabo's correction term and lens surgery* Math. Proc. Cambridge Philos. Soc. 146 (2009) , no.1 119–134

[8] M. Tange, *Lens spaces given from L-space homology 3-spheres* Experiment. Math. 18, no 3 (2009) 285–301.

[9] J. Rasmussen, *Lens space surgeries and L-space homology spheres* arXiv:0710.2531v1

Motoo Tange
Research Institute for Mathematical Sciences,
Kyoto University,
Kyoto 606-8502, Japan.
tange@kurims.kyoto-u.ac.jp